

## 26.4 The Role of Seed Plants

*By the end of this section, you will be able to do the following:*

- Explain how angiosperm diversity is due, in part, to multiple complex interactions with animals
- Describe ways in which pollination occurs
- Discuss the roles that plants play in ecosystems and how deforestation threatens plant biodiversity

Without seed plants, life as we know it would not be possible. Plants play a key role in the maintenance of terrestrial ecosystems through the stabilization of soils, cycling of carbon, and climate moderation. Large tropical forests release oxygen and act as carbon dioxide “sinks.” Seed plants provide shelter to many life forms, as well as food for herbivores, thereby indirectly feeding carnivores. Plant secondary metabolites are used for medicinal purposes and industrial production. Virtually all animal life is dependent on plants for survival.

### Animals and Plants: Herbivory

Coevolution of flowering plants and insects is a hypothesis that has received much attention and support, especially because both angiosperms and insects diversified at about the same time in the middle Mesozoic. Many authors have attributed the diversity of plants and insects to both pollination and **herbivory**, or the consumption of plants by insects and other animals. Herbivory is believed to have been as much a driving force as pollination. Coevolution of herbivores and plant defenses is easily and commonly observed in nature. Unlike animals, most plants cannot outrun predators or use mimicry to hide from hungry animals (although mimicry has been used to entice pollinators). A sort of arms race exists between plants and herbivores. To “combat” herbivores, some plant seeds—such as acorn and unripened persimmon—are high in alkaloids and therefore unsavory to some animals. Other plants are protected by bark, although some animals developed specialized mouth pieces to tear and chew vegetal material. Spines and thorns (Figure 26.20) deter most animals, except for mammals with thick fur, and some birds have specialized beaks to get past such defenses.



**Figure 26.20** Plant defenses. (a) Spines and (b) thorns are examples of plant defenses. (credit a: modification of work by Jon Sullivan; credit b: modification of work by I. Sáček, Sr.)

Herbivory has been exploited by seed plants for their own benefit. The dispersal of fruits by herbivorous animals is a striking example of mutualistic relationships. The plant offers to the herbivore a nutritious source of food in return for spreading the plant’s genetic material to a wider area.

An extreme example of coevolution (discovered by Daniel Janzen) between an animal and a plant is exemplified by Mexican acacia trees and their attendant acacia ants *Pseudomyrmex* spp. (this is termed *myrmecophytism*). The trees support the ants with shelter and food: The ants nest in the hollows of large thorns produced by the tree and feed on sugary secretions produced at the ends of the leaves. The sugar pellets also help to keep the ants from interfering with insect pollinators. In return, ants discourage herbivores, both invertebrates and vertebrates, by stinging and attacking leaf-eaters and insects ovipositing on the plants. The ants also help to remove potential plant pathogens, such as fungal growths. Another case of insect-plant coevolution is found in bracken fern (*Pteridium aquilinum*), whose subspecies are found throughout the world. Bracken ferns produce a number of “secondary plant compounds” in their adult fronds that serve as defensive compounds against nonadapted insect attack (these compounds include cyanogenic glucosides, tannins, and phenolics). However, during the “fiddlehead” or crozier stage, bracken secretes nutritious sugary and proteinaceous compounds from special “nectaries” that attract ants and even species of jumping spiders, all of which defend the plant’s croziers until they are fully unfolded. These opportunistic groups of protective arthropods greatly reduce the damage that otherwise would occur during the early stages of growth.


## Animals and Plants: Pollination

Flowers pollinated by wind are usually small, feathery, and visually inconspicuous. *Grasses* are a successful group of flowering plants that are wind pollinated. They produce large amounts of powdery pollen carried over large distances by the wind. Some large trees such as oaks, maples, and birches are also wind pollinated.

### LINK TO LEARNING

Explore this [website \(http://openstax.org/l/pollinators2\)](http://openstax.org/l/pollinators2) for additional information on pollinators.

More than 80 percent of angiosperms depend on animals for **pollination** (technically the transfer of pollen from the anther to the stigma). Consequently, plants have developed many adaptations to attract pollinators. With over 200,000 different plants dependent on animal pollination, the plant needs to advertise to its pollinators with some specificity. The specificity of specialized plant structures that target animals can be very surprising. It is possible, for example, to determine the general type of pollinators favored by a plant by observing the flower's physical characteristics. Many bird or insect-pollinated flowers secrete nectar, which is a sugary liquid. They also produce both fertile pollen, for reproduction, and sterile pollen rich in nutrients for birds and insects. Many butterflies and bees can detect ultraviolet light, and flowers that attract these pollinators usually display a pattern of ultraviolet reflectance that helps them quickly locate the flower's center. In this manner, pollinating insects collect nectar while at the same time are dusted with pollen ([Figure 26.21](#)). Large, red flowers with little smell and a long funnel shape are preferred by hummingbirds, who have good color perception, a poor sense of smell, and need a strong perch. White flowers that open at night attract moths. Other animals—such as bats, lemurs, and lizards—can also act as pollinating agents. Any disruption to these interactions, such as the disappearance of bees, for example as a consequence of colony collapse disorders, can lead to disaster for agricultural industries that depend heavily on pollinated crops.



**Figure 26.21** Pollination. As a bee collects nectar from a flower, it is dusted by pollen, which it then disperses to other flowers. (credit: John Severns)

### SCIENTIFIC METHOD CONNECTION

#### Testing Attraction of Flies by Rotting Flesh Smell

**Question:** Will flowers that offer cues to bees attract carrion flies if sprayed with compounds that smell like rotten flesh?

**Background:** Visitation of flowers by pollinating flies is a function mostly of smell. Flies are attracted by rotting flesh and carrions. The putrid odor seems to be the major attractant. The polyamines putrescine and cadaverine, which are the products of protein breakdown after animal death, are the source of the pungent smell of decaying meat. Some plants strategically attract flies by synthesizing polyamines similar to those generated by decaying flesh and thereby attract carrion flies.

Flies seek out dead animals because they normally lay their eggs on them and their maggots feed on the decaying flesh. Interestingly, time of death can be determined by a forensic entomologist based on the stages and type of maggots recovered from cadavers.

**Hypothesis:** Because flies are drawn to other organisms based on smell and not sight, a flower that is normally attractive to bees because of its colors will attract flies if it is sprayed with polyamines similar to those generated by decaying flesh.

**Test the hypothesis:**

1. Select flowers usually pollinated by bees. White petunia may be a good choice.
2. Divide the flowers into two groups, and while wearing eye protection and gloves, spray one group with a solution of either putrescine or cadaverine. (Putrescine dihydrochloride is typically available in 98 percent concentration; this can be diluted to approximately 50 percent for this experiment.)
3. Place the flowers in a location where flies are present, keeping the sprayed and unsprayed flowers separated.
4. Observe the movement of the flies for one hour. Record the number of visits to the flowers using a table similar to [Table 26.2](#). Given the rapid movement of flies, it may be beneficial to use a video camera to record the fly–flower interaction. Replay the video in slow motion to obtain an accurate record of the number of fly visits to the flowers.
5. Repeat the experiment four more times with the same species of flower, but using different specimens.
6. Repeat the entire experiment with a different type of flower that is normally pollinated by bees.

Results of Number of Visits by Flies to Sprayed and Control/Unsprayed Flowers

Trial #	Sprayed Flowers	Unsprayed Flowers
1		
2		
3		
4		
5		

**Table 26.2**

**Analyze your data:** Review the data you have recorded. Average the number of visits that flies made to sprayed flowers over the course of the five trials (on the first flower type) and compare and contrast them to the average number of visits that flies made to the unsprayed/control flowers. Can you draw any conclusions regarding the attraction of the flies to the sprayed flowers?

For the second flower type used, average the number of visits that flies made to sprayed flowers over the course of the five trials and compare and contrast them to the average number of visits that flies made to the unsprayed/control flowers. Can you draw any conclusions regarding the attraction of the flies to the sprayed flowers?

Compare and contrast the average number of visits that flies made to the two flower types. Can you draw any conclusions about whether the appearance of the flower had any impact on the attraction of flies? Did smell override any appearance differences, or were the flies attracted to one flower type more than another?

**Form a conclusion:** Do the results support the hypothesis? If not, how can your observations be explained?

## The Importance of Seed Plants in Human Life

Seed plants are the foundation of human diets across the world ([Figure 26.22](#)). Many societies eat almost exclusively vegetarian fare and depend solely on seed plants for their nutritional needs. A few crops (rice, wheat, and potatoes) dominate the agricultural landscape. Many crops were developed during the agricultural revolution, when human societies made the transition from nomadic hunter–gatherers to horticulture and agriculture. Cereals, rich in carbohydrates, provide the staple of many human diets. Beans and nuts supply proteins. Fats are derived from crushed seeds, as is the case for peanut and rapeseed (canola) oils, or fruits such as olives. Animal husbandry also consumes large quantities of crop plants.

Staple crops are not the only food derived from seed plants. Various fruits and vegetables provide nutrient macromolecules,

vitamins, minerals, and fiber. Sugar, to sweeten dishes, is produced from the monocot sugarcane and the eudicot sugar beet. Drinks are made from infusions of tea leaves, chamomile flowers, crushed coffee beans, or powdered cocoa beans. Spices come from many different plant parts: saffron and cloves are stamens and buds, black pepper and vanilla are seeds, the bark of a bush in the *Laurales* family supplies cinnamon, and the herbs that flavor many dishes come from dried leaves and fruit, such as the pungent red chili pepper. The volatile oils of a number of flowers and bark provide the scent of perfumes.

Additionally, no discussion of seed plant contribution to human diet would be complete without the mention of alcohol. Fermentation of plant-derived sugars and starches is used to produce alcoholic beverages in all societies. In some cases, the beverages are derived from the fermentation of sugars from fruit, as with wines and, in other cases, from the fermentation of carbohydrates derived from seeds, as with beers. The sharing of foods and beverages also contributes to human social ritual.

Seed plants have many other uses, including providing wood as a source of timber for construction, fuel, and material to build furniture. Most paper is derived from the pulp of coniferous trees. Fibers of seed plants such as cotton, flax, and hemp are woven into cloth. Textile dyes, such as indigo, were mostly of plant origin until the advent of synthetic chemical dyes.

Lastly, it is more difficult to quantify the benefits of ornamental seed plants. These grace private and public spaces, adding beauty and serenity to human lives and inspiring painters and poets alike.



**Figure 26.22** Human uses of plants. Humans rely on plants for a variety of reasons. (a) Cacao beans were introduced to Europe from the New World, where they were used by Mesoamerican civilizations. Combined with sugar, another plant product, chocolate is a popular food. (b) Flowers like the tulip are cultivated for their beauty. (c) Quinine, extracted from cinchona trees, is used to treat malaria, to reduce fever, and to alleviate pain. (d) This violin is made of wood. (credit a: modification of work by "Everjean"/Flickr; credit b: modification of work by Rosendahl; credit c: modification of work by Franz Eugen Köhler)

The medicinal properties of plants have been known to human societies since ancient times. There are references to the use of plants' curative properties in Egyptian, Babylonian, and Chinese writings from 5,000 years ago. Many modern synthetic therapeutic drugs are derived or synthesized *de novo* from plant secondary metabolites. It is important to note that the same plant extract can be a therapeutic remedy at low concentrations, become an addictive drug at higher doses, and can potentially kill at high concentrations. [Table 26.3](#) presents a few drugs, their plants of origin, and their medicinal applications.

Plant Origin of Medicinal Compounds and Medical Applications

Plant	Compound	Application
Deadly nightshade ( <i>Atropa belladonna</i> )	Atropine	Dilate eye pupils for eye exams
Foxglove ( <i>Digitalis purpurea</i> )	Digitalis	Heart disease, stimulates heart beat
Yam ( <i>Dioscorea</i> spp.)	Steroids	Steroid hormones: contraceptive pill and cortisone
Ephedra ( <i>Ephedra</i> spp.)	Ephedrine	Decongestant and bronchiole dilator
Pacific yew ( <i>Taxus brevifolia</i> )	Taxol	Cancer chemotherapy; inhibits mitosis
Opium poppy ( <i>Papaver somniferum</i> )	Opioids	Analgesic (reduces pain without loss of consciousness) and narcotic (reduces pain with drowsiness and loss of consciousness) in higher doses
Quinine tree ( <i>Cinchona</i> spp.)	Quinine	Antipyretic (lowers body temperature) and antimalarial
Willow ( <i>Salix</i> spp.)	Salicylic acid (aspirin)	Analgesic and antipyretic

Table 26.3



## CAREER CONNECTION

### Ethnobotanist

The relatively new field of ethnobotany studies the interaction between a particular culture and the plants native to the region. Seed plants have a large influence on day-to-day human life. Not only are plants the major source of food and medicine, they also influence many other aspects of society, from clothing to industry. The medicinal properties of plants were recognized early on in human cultures. From the mid-1900s, synthetic chemicals began to supplant plant-based remedies.

Pharmacognosy is the branch of pharmacology that focuses on medicines derived from natural sources. With massive globalization and industrialization, it is possible that much human knowledge of plants and their medicinal purposes will disappear with the cultures that fostered them. This is where ethnobotanists come in. To learn about and understand the use of plants in a particular culture, an ethnobotanist must bring in knowledge of plant life and an understanding and appreciation of diverse cultures and traditions. The Amazon forest is home to an incredible diversity of vegetation and is considered an untapped resource of medicinal plants; yet, both the ecosystem and its indigenous cultures are threatened with extinction.

To become an ethnobotanist, a person must acquire a broad knowledge of plant biology, ecology, and sociology. Not only are the



plant specimens studied and collected, but also the stories, recipes, and traditions that are linked to them. For ethnobotanists, plants are not viewed solely as biological organisms to be studied in a laboratory, but as an integral part of human culture. The convergence of molecular biology, anthropology, and ecology make the field of ethnobotany a truly multidisciplinary science.

---

## Biodiversity of Plants

Biodiversity ensures a resource for new food crops and medicines. Plant life balances ecosystems, protects watersheds, mitigates erosion, moderates our climate, and provides shelter for many animal species. Threats to plant diversity, however, come from many sources. The explosion of the human population, especially in tropical countries where birth rates are highest and economic development is in full swing, is leading to devastating human encroachment into forested areas. To feed the growing population, humans need to obtain arable land, so there has been and continues to be massive clearing of trees. The need for more energy to power larger cities and economic growth therein leads to the construction of dams, the consequent flooding of ecosystems, and increased emissions of pollutants. Other threats to tropical forests come from poachers, who log trees for their precious wood. Ebony and Brazilian rosewood, both on the endangered list, are examples of tree species driven almost to extinction by indiscriminate logging. This unfortunate practice continues unabated today largely due to lack of population control and political willpower.

The number of plant species becoming extinct is increasing at an alarming rate. Because ecosystems are in a delicate balance, and seed plants maintain close symbiotic relationships with animals—whether predators or pollinators—the disappearance of a single plant can lead to the extinction of connected animal species. A real and pressing issue is that many plant species have not yet been catalogued, and so their place in the ecosystem is unknown. These unknown species are threatened by logging, habitat destruction, and loss of pollinators. They may become extinct before we have the chance to begin to understand the possible impacts from their disappearance. Efforts to preserve biodiversity take several lines of action, from preserving heirloom seeds to barcoding species. **Heirloom seeds** come from plants that were traditionally grown in human populations, as opposed to the seeds used for large-scale agricultural production. **Barcoding** is a technique in which one or more short gene sequences, taken from a well-characterized portion of DNA found in most genomes, are used to identify a species through DNA analysis.

## KEY TERMS

**anther** sac-like structure at the tip of the stamen in which pollen grains are produced

**Anthophyta** phylum to which angiosperms belong

**barcoding** molecular biology technique in which one or more short gene sequences taken from a well-characterized portion of the genome is used to identify a species

**basal angiosperms** a group of plants that probably branched off before the separation of monocots and eudicots

**calyx** whorl of sepals

**carpel** single unit of the pistil

**conifer** dominant phylum of gymnosperms with the greatest variety of trees

**corolla** collection of petals

**cotyledon** primitive leaf that develops in the zygote; monocots have one cotyledon, and dicots have two cotyledons

**crop** cultivated plant

**cycad** gymnosperm that grows in tropical climates and resembles a palm tree; member of the phylum Cycadophyta

**dicot** (also, eudicot) related group of angiosperms whose embryos possess two cotyledons

**dioecious** describes a species in which the male and female reproductive organs are carried on separate specimens

**filament** thin stalk that links the anther to the base of the flower

**flower** branches specialized for reproduction found in some seed-bearing plants, containing either specialized male or female organs or both male and female organs

**fruit** thickened tissue derived from ovary wall that protects the embryo after fertilization and facilitates seed dispersal

**ginkgophyte** gymnosperm with one extant species, the *Ginkgo biloba*: a tree with fan-shaped leaves

**gnetophyte** gymnosperm shrub with varied morphological features that produces vessel elements in its woody tissues; the phylum includes the genera *Ephedra*, *Gnetum*, and *Welwitschia*

**gymnosperm** seed plant with naked seeds (seeds exposed on modified leaves or in cones)

**gynoecium** (also, carpel) structure that constitutes the female reproductive organ

**heirloom seed** seed from a plant that was grown historically, but has not been used in modern agriculture on a large scale

**herbaceous** grass-like plant noticeable by the absence of woody tissue

**herbivory** consumption of plants by insects and other animals

**integument** layer of sporophyte tissue that surrounds the megasporangium, and later, the embryo

**megasporeocyte** megaspore mother cell; larger spore that germinates into a female gametophyte in a heterosporous plant

**microsporeocyte** smaller spore that produces a male gametophyte in a heterosporous plant

**monocot** related group of angiosperms that produce embryos with one cotyledon and pollen with a single ridge

**monoecious** describes a species in which the male and female reproductive organs are on the same plant

**nectar** liquid rich in sugars produced by flowers to attract animal pollinators

**ovary** chamber that contains and protects the ovule or female megasporangium

**ovulate cone** cone containing two ovules per scale

**ovule** female gametophyte

**perianth** part of the plant consisting of the calyx (sepals) and corolla (petals)

**petal** modified leaf interior to the sepals; colorful petals attract animal pollinators

**pistil** fused group of carpels

**pollen grain** structure containing the male gametophyte of the plant

**pollen tube** extension from the pollen grain that delivers sperm to the egg cell

**pollination** transfer of pollen from the anther to the stigma

**progymnosperm** transitional group of plants that resembled conifers because they produced wood, yet still reproduced like ferns

**seed** structure containing the embryo, storage tissue, and protective coat

**sepal** modified leaf that encloses the bud; outermost structure of a flower

**spermatophyte** seed plant; from the Greek *sperm* (seed) and *phyte* (plant)

**stamen** structure that contains the male reproductive organs

**stigma** uppermost structure of the carpel where pollen is deposited

**strobilus** plant structure with a tight arrangement of sporophylls around a central stalk, as seen in cones or flowers; the male strobilus produces pollen, and the female strobilus produces eggs

**style** long, thin structure that links the stigma to the ovary

## CHAPTER SUMMARY

### 26.1 Evolution of Seed Plants

Seed plants appeared about 350 million years ago, during the Carboniferous period. Two major innovations were seeds and pollen. Seeds protect the embryo from desiccation and provide it with a store of nutrients to support the early growth of the sporophyte. Seeds are also equipped to delay germination until growth conditions are optimal. Pollen allows seed plants to reproduce in the absence of water. The gametophytes of seed plants shrank, while the sporophytes became prominent structures and the diploid stage became the longest phase of the life cycle.

In the gymnosperms, which appeared during the drier Permian period and became the dominant group during the Triassic, pollen was dispersed by wind, and their naked seeds developed in the sporophylls of a strobilus. Angiosperms bear both flowers and fruit. Flowers expand the possibilities for pollination, especially by insects, who have coevolved with the flowering plants. Fruits offer additional protection to the embryo during its development, and also assist with seed dispersal. Angiosperms appeared during the Mesozoic era and have become the dominant plant life in terrestrial habitats.

### 26.2 Gymnosperms

Gymnosperms are heterosporous seed plants that produce naked seeds. They appeared in the Paleozoic period and were the dominant plant life during the Mesozoic. Modern-day gymnosperms belong to four phyla. The largest phylum, Coniferophyta, is represented by conifers, the predominant plants at high altitude and latitude. Cycads (phylum Cycadophyta) resemble palm trees and grow in tropical climates. Ginkgophyta is represented today by a single species, *Ginkgo biloba*. The last phylum, Gnetophyta, is a diverse group of plants that produce vessel elements in their wood.

### 26.3 Angiosperms

Angiosperms are the dominant form of plant life in most terrestrial ecosystems, comprising about 90 percent of all

plant species. Most crops and ornamental plants are angiosperms. Their success comes from two innovative structures that protect reproduction from variability in the environment: the flower and the fruit. Flowers were derived from modified leaves; their color and fragrance encourages species-specific pollination. The main parts of a flower are the sepals and petals, which protect the reproductive parts: the stamens and the carpels. The stamens produce the male gametes in pollen grains. The carpels contain the female gametes (the eggs inside the ovules), which are within the ovary of a carpel. The walls of the ovary thicken after fertilization, ripening into fruit that ensures dispersal by wind, water, or animals.

The angiosperm life cycle is dominated by the sporophyte stage. Double fertilization is an event unique to angiosperms. One sperm in the pollen fertilizes the egg, forming a diploid zygote, while the other combines with the two polar nuclei, forming a triploid cell that develops into a food storage tissue called the endosperm. Flowering plants are divided into two main groups, the monocots and eudicots, according to the number of cotyledons in the seedlings. Basal angiosperms belong to an older lineage than monocots and eudicots.

### 26.4 The Role of Seed Plants

Angiosperm diversity is due in part to multiple interactions with animals. Herbivory has favored the development of defense mechanisms in plants, and avoidance of those defense mechanisms in animals. Conversely, seed dispersal can be aided by animals that eat plant fruits. Pollination (the transfer of pollen to a carpel) is mainly carried out by wind and animals, and angiosperm fruits and seeds have evolved numerous adaptations to capture the wind or attract specific classes of animals.

Plants play a key role in ecosystems. They are a source of food and medicinal compounds, and provide raw materials for many industries. Rapid deforestation and industrialization, however, threaten plant biodiversity. In turn, this threatens the ecosystem.

## VISUAL CONNECTION QUESTIONS

- Figure 26.8** At what stage does the diploid zygote form?
  - when the female cone begins to bud from the tree
  - at fertilization
  - when the seeds drop from the tree
  - when the pollen tube begins to grow
- Figure 26.15** If a flower lacked a megasporangium, what type of gamete would not form? If the flower lacked a microsporangium, what type of gamete would not form?



## REVIEW QUESTIONS

3. Seed plants are \_\_\_\_\_.
  - a. all homosporous
  - b. mostly homosporous with some heterosporous
  - c. mostly heterosporous with some homosporous
  - d. all heterosporous
4. Besides the seed, what other major structure diminishes a plant's reliance on water for reproduction?
  - a. flower
  - b. fruit
  - c. pollen
  - d. spore
5. In which of the following geological periods would gymnosperms dominate the landscape?
  - a. Carboniferous
  - b. Permian
  - c. Triassic
  - d. Eocene (present)
6. Which of the following structures widens the geographic range of a species and is an agent of dispersal?
  - a. seed
  - b. flower
  - c. leaf
  - d. root
7. Which of the following traits characterizes gymnosperms?
  - a. The plants carry exposed seeds on modified leaves.
  - b. Reproductive structures are located in a flower.
  - c. After fertilization, the ovary thickens and forms a fruit.
  - d. The gametophyte is the longest phase of the life cycle.
8. Megaspores will eventually produce which of the following?
  - a. pollen grain
  - b. sporophytes
  - c. male gametophytes
  - d. female gametophytes
9. What is the ploidy of the following structures: gametophyte, seed, spore, sporophyte?
  - a.  $1n$ ,  $1n$ ,  $2n$ ,  $2n$
  - b.  $1n$ ,  $2n$ ,  $1n$ ,  $2n$
  - c.  $2n$ ,  $1n$ ,  $2n$ ,  $1n$
  - d.  $2n$ ,  $2n$ ,  $1n$ ,  $1n$
10. In the northern forests of Siberia, a tall tree is most likely a:
  - a. conifer
  - b. cycad
  - c. *Ginkgo biloba*
  - d. gnetophyte
11. Which of the following structures in a flower is not directly involved in reproduction?
  - a. the style
  - b. the stamen
  - c. the sepal
  - d. the anther
12. Pollen grains develop in which structure?
  - a. the anther
  - b. the stigma
  - c. the filament
  - d. the carpel
13. In the course of double fertilization, one sperm cell fuses with the egg and the second one fuses with \_\_\_\_\_.
  - a. the synergids
  - b. the polar nuclei of the center cell
  - c. the egg as well
  - d. the antipodal cells
14. Corn develops from a seedling with a single cotyledon, displays parallel veins on its leaves, and produces monosulcate pollen. It is most likely:
  - a. a gymnosperm
  - b. a monocot
  - c. a eudicot
  - d. a basal angiosperm
15. Which of the following plant structures is not a defense against herbivory?
  - a. thorns
  - b. spines
  - c. nectar
  - d. alkaloids
16. White and sweet-smelling flowers with abundant nectar are probably pollinated by
  - a. bees and butterflies
  - b. flies
  - c. birds
  - d. wind

17. Abundant and powdery pollen produced by small, indistinct flowers is probably transported by:
  - a. bees and butterflies
  - b. flies
  - c. birds
  - d. wind
18. Plants are a source of \_\_\_\_\_.
  - a. food
  - b. fuel
  - c. medicine
  - d. all of the above

## CRITICAL THINKING QUESTIONS

19. The Cretaceous Period was marked by the increase in number and variety of angiosperms. Insects also diversified enormously during the same period. Can you propose the reason or reasons that could foster coevolution?
20. What role did the adaptations of seed and pollen play in the development and expansion of seed plants?
21. The Mediterranean landscape along the sea shore is dotted with pines and cypresses. The weather is not cold, and the trees grow at sea level. What evolutionary adaptation of conifers makes them suitable to the Mediterranean climate?
22. What are the four modern-day phyla of gymnosperms?
23. Some cycads are considered endangered species and their trade is severely restricted. Customs officials stop suspected smugglers who claim that the plants in their possession are palm trees, not cycads. How would a botanist distinguish between the two types of plants?
24. What are the two structures that allow angiosperms to be the dominant form of plant life in most terrestrial ecosystems?
25. Biosynthesis of nectar and nutrient-rich pollen is energetically very expensive for a plant. Yet, plants funnel large amounts of energy into animal pollination. What are the evolutionary advantages that offset the cost of attracting animal pollinators?
26. What is biodiversity and why is it important to an ecosystem?

