CHAPTER 21 The Quantum Nature of Light



Figure 21.1 In Lewis Carroll's classic text Alice's Adventures in Wonderland, Alice follows a rabbit down a hole into a land of curiosity. While many of her interactions in Wonderland are of surprising consequence, they follow a certain inherent logic. (credit: modification of work by John Tenniel, Wikimedia Commons)

Chapter Outline

21.1 Planck and Quantum Nature of Light

21.2 Einstein and the Photoelectric Effect

21.3 The Dual Nature of Light

INTRODUCTION At first glance, the quantum nature of light can be a strange and bewildering concept. Between light acting as discrete chunks, massless particles providing momenta, and fundamental particles behaving like waves, it may often seem like something out of Alice in Wonderland.

For many, the study of this branch of physics can be as enthralling as Lewis Carroll's classic novel. Recalling the works of legendary characters and brilliant scientists such as Einstein, Planck, and Compton, the study of light's quantum nature will provide you an interesting tale of how a clever interpretation of some small details led to the most important discoveries of the past 150 years. From the electronics revolution of the twentieth century to our future progress in solar energy and space exploration, the quantum nature of light should yield a rabbit hole of curious consequence, within which lie some of the most fascinating truths of our time.

21.1 Planck and Quantum Nature of Light

Section Learning Objectives

By the end of this section, you will be able to do the following:

- Describe blackbody radiation
- Define quantum states and their relationship to modern physics
- Calculate the quantum energy of lights
- Explain how photon energies vary across divisions of the electromagnetic spectrum

Section Key Terms

blackbody quantized quantum ultraviolet catastrophe

Blackbodies

Our first story of curious significance begins with a T-shirt. You are likely aware that wearing a tight black T-shirt outside on a hot day provides a significantly less comfortable experience than wearing a white shirt. Black shirts, as well as all other black objects, will absorb and re-emit a significantly greater amount of radiation from the sun. This shirt is a good approximation of what is called a **blackbody**.

A perfect blackbody is one that absorbs and re-emits all radiated energy that is incident upon it. Imagine wearing a tight shirt that did this! This phenomenon is often modeled with quite a different scenario. Imagine carving a small hole in an oven that can be heated to very high temperatures. As the temperature of this container gets hotter and hotter, the radiation out of this dark hole would increase as well, re-emitting all energy provided it by the increased temperature. The hole may even begin to glow in different colors as the temperature is increased. Like a burner on your stove, the hole would glow red, then orange, then blue, as the temperature is increased. In time, the hole would continue to glow but the light would be invisible to our eyes. This container is a good model of a perfect blackbody.

It is the analysis of blackbodies that led to one of the most consequential discoveries of the twentieth century. Take a moment to carefully examine Figure 21.2. What relationships exist? What trends can you see? The more time you spend interpreting this figure, the closer you will be to understanding quantum physics!



Figure 21.2 Graphs of blackbody radiation (from an ideal radiator) at three different radiator temperatures. The intensity or rate of radiation emission increases dramatically with temperature, and the peak of the spectrum shifts toward the visible and ultraviolet parts of the spectrum. The shape of the spectrum cannot be described with classical physics.

TIPS FOR SUCCESS

When encountering a new graph, it is best to try to interpret the graph before you read about it. Doing this will make the following text more meaningful and will help to remind yourself of some of the key concepts within the section.

Understanding Blackbody Graphs

<u>Figure 21.2</u> is a plot of radiation intensity against radiated wavelength. In other words, it shows how the intensity of radiated light changes when a blackbody is heated to a particular temperature.

It may help to just follow the bottom-most red line labeled 3,000 K, red hot. The graph shows that when a blackbody acquires a temperature of 3,000 K, it radiates energy across the electromagnetic spectrum. However, the energy is most intensely emitted at a wavelength of approximately 1000 nm. This is in the infrared portion of the electromagnetic spectrum. While a body at this temperature would appear *red-hot* to our eyes, it would truly appear 'infrared-hot' if we were able to see the entire spectrum.

A few other important notes regarding <u>Figure 21.2</u>:

- As temperature increases, the total amount of energy radiated increases. This is shown by examining the area underneath each line.
- Regardless of temperature, all red lines on the graph undergo a consistent pattern. While electromagnetic radiation is emitted throughout the spectrum, the intensity of this radiation peaks at one particular wavelength.
- As the temperature changes, the wavelength of greatest radiation intensity changes. At 4,000 K, the radiation is most intense in the yellow-green portion of the spectrum. At 6,000 K, the blackbody would radiate *white hot*, due to intense radiation throughout the visible portion of the electromagnetic spectrum. Remember that white light is the emission of all visible colors simultaneously.
- As the temperature increases, the frequency of light providing the greatest intensity increases as well. Recall the equation $v = f\lambda$. Because the speed of light is constant, frequency and wavelength are inversely related. This is verified by the leftward movement of the three red lines as temperature is increased.

While in science it is important to categorize observations, theorizing as to why the observations exist is crucial to scientific advancement. Why doesn't a blackbody emit radiation evenly across all wavelengths? Why does the temperature of the body change the peak wavelength that is radiated? Why does an increase in temperature cause the peak wavelength emitted to decrease? It is questions like these that drove significant research at the turn of the twentieth century. And within the context of these questions, Max Planck discovered something of tremendous importance.

Planck's Revolution

The prevailing theory at the time of Max Planck's discovery was that intensity and frequency were related by the equation $I = \frac{2kT}{\lambda^2}$. This equation, derived from classical physics and using wave phenomena, infers that as wavelength increases, the intensity of energy provided will decrease with an inverse-squared relationship. This relationship is graphed in Figure 21.3 and shows a troubling trend. For starters, it should be apparent that the graph from this equation does not match the blackbody graphs found experimentally. Additionally, it shows that for an object of any temperature, there should be an infinite amount of energy quickly emitted in the shortest wavelengths. When theory and experimental results clash, it is important to re-evaluate

both models. The disconnect between theory and reality was termed the ultraviolet catastrophe.



Figure 21.3 The graph above shows the true spectral measurements by a blackbody against those predicted by the classical theory at the time. The discord between the predicted classical theory line and the actual results is known as the ultraviolet catastrophe.

Due to concerns over the ultraviolet catastrophe, Max Planck began to question whether another factor impacted the relationship between intensity and wavelength. This factor, he posited, should affect the probability that short wavelength light would be emitted. Should this factor reduce the probability of short wavelength light, it would cause the radiance curve to not progress infinitely as in the classical theory, but would instead cause the curve to precipitate back downward as is shown in the 5,000 K, 4,000 K, and 3,000 K temperature lines of the graph in Figure 21.3. Planck noted that this factor, whatever it may be, must also be dependent on temperature, as the intensity decreases at lower and lower wavelengths as the temperature increases.

The determination of this *probability factor* was a groundbreaking discovery in physics, yielding insight not just into light but also into energy and matter itself. It would be the basis for Planck's 1918 Nobel Prize in Physics and would result in the transition of physics from classical to modern understanding. In an attempt to determine the cause of the *probability factor*, Max Planck constructed a new theory. This theory, which created the branch of physics called quantum mechanics, speculated that the energy radiated by the blackbody could exist only in specific numerical, or **quantum**, states. This theory is described by the equation E = nhf, where *n* is any nonnegative integer (0, 1, 2, 3, ...) and *h* is Planck's constant, given by $h = 6.626 \times 10^{-34} \text{ J} \cdot \text{s}$, and *f* is frequency.

Through this equation, Planck's probability factor can be more clearly understood. Each frequency of light provides a specific quantized amount of energy. Low frequency light, associated with longer wavelengths would provide a smaller amount of energy, while high frequency light, associated with shorter wavelengths, would provide a larger amount of energy. For specified temperatures with specific total energies, it makes sense that more low frequency light would be radiated than high frequency light. To a degree, the relationship is like pouring coins through a funnel. More of the smaller pennies would be able to pass through the funnel than the larger quarters. In other words, because the value of the coin is somewhat related to the size of the coin, the probability of a quarter passing through the funnel is reduced!

Furthermore, an increase in temperature would signify the presence of higher energy. As a result, the greater amount of total blackbody energy would allow for more of the high frequency, short wavelength, energies to be radiated. This permits the peak of the blackbody curve to drift leftward as the temperature increases, as it does from the 3,000 K to 4,000 K to 5,000 K values. Furthering our coin analogy, consider a wider funnel. This funnel would permit more quarters to pass through and allow for a reduction in concern about the *probability factor*.

In summary, it is the interplay between the predicted classical model and the quantum probability that creates the curve depicted in <u>Figure 21.3</u>. Just as quarters have a higher currency denomination than pennies, higher frequencies come with larger

21.1

amounts of energy. However, just as the probability of a quarter passing through a fixed diameter funnel is reduced, so is the probability of a high frequency light existing in a fixed temperature object. As is often the case in physics, it is the balancing of multiple incredible ideas that finally allows for better understanding.

Quantization

It may be helpful at this point to further consider the idea of quantum states. Atoms, molecules, and fundamental electron and proton charges are all examples of physical entities that are **quantized**—that is, they appear only in certain discrete values and do not have every conceivable value. On the macroscopic scale, this is not a revolutionary concept. A standing wave on a string allows only particular harmonics described by integers. Going up and down a hill using discrete stair steps causes your potential energy to take on discrete values as you move from step to step. Furthermore, we cannot have a fraction of an atom, or part of an electron's charge, or 14.33 cents. Rather, everything is built of integral multiples of these substructures.

That said, to discover quantum states within a phenomenon that science had always considered continuous would certainly be surprising. When Max Planck was able to use quantization to correctly describe the experimentally known shape of the blackbody spectrum, it was the first indication that energy was quantized on a small scale as well. This discovery earned Planck the Nobel Prize in Physics in 1918 and was such a revolutionary departure from classical physics that Planck himself was reluctant to accept his own idea. The general acceptance of Planck's energy quantization was greatly enhanced by Einstein's explanation of the photoelectric effect (discussed in the next section), which took energy quantization a step further.



Figure 21.4 The German physicist Max Planck had a major influence on the early development of quantum mechanics, being the first to recognize that energy is sometimes quantized. Planck also made important contributions to special relativity and classical physics. (credit: Library of Congress, Prints and Photographs Division, Wikimedia Commons)

How Many Photons per Second Does a Typical Light Bulb Produce?

Assuming that 10 percent of a 100-W light bulb's energy output is in the visible range (typical for incandescent bulbs) with an average wavelength of 580 nm, calculate the number of visible photons emitted per second.

Strategy

The number of visible photons per second is directly related to the amount of energy emitted each second, also known as the bulb's power. By determining the bulb's power, the energy emitted each second can be found. Since the power is given in watts, which is joules per second, the energy will be in joules. By comparing this to the amount of energy associated with each photon, the number of photons emitted each second can be determined.

Solution

The power in visible light production is 10.0 percent of 100 W, or 10.0 J/s. The energy of the average visible photon is found by substituting the given average wavelength into the formula

$$E = nhf = \frac{nhc}{\lambda}$$

By rearranging the above formula to determine energy per photon, this produces

$$E/n = \frac{(6.63 \times 10^{-34} \text{J} \cdot \text{s})(3.00 \times 10^8 \text{ m/s})}{580 \times 10^{-9} \text{m}} = 3.43 \times 10^{-19} \text{J/photon.}$$

The number of visible photons per second is thus

$$\frac{\text{photons}}{\text{sec}} = \frac{10.0 \text{J/s}}{3.43 \times 10^{-19} \text{J/photon}} = 2.92 \times 10^{19} \text{photons/s}.$$

Discussion

This incredible number of photons per second is verification that individual photons are insignificant in ordinary human experience. However, it is also a verification of our everyday experience—on the macroscopic scale, photons are so small that quantization becomes essentially continuous.



How does Photon Energy Change with Various Portions of the EM Spectrum?

Refer to the Graphs of Blackbody Radiation shown in the first figure in this section. Compare the energy necessary to radiate one photon of infrared light and one photon of visible light.

Strategy

To determine the energy radiated, it is necessary to use the equation E = nhf. It is also necessary to find a representative frequency for infrared light and visible light.

Solution

According to the first figure in this section, one representative wavelength for infrared light is 2000 nm (2.000 \times 10⁻⁶ m). The associated frequency of an infrared light is

$$f = \frac{c}{\lambda} = \frac{3.00 \times 10^8 \text{ m/s}}{2.000 \times 10^{-6} \text{ m}} = 1.50 \times 10^{14} \text{ Hz.}$$
 21.2

Using the equation E = nhf, the energy associated with one photon of representative infrared light is

$$\frac{E}{n} = h \cdot f = (6.63 \times 10^{-34} \text{J} \cdot \text{s}) (1.50 \times 10^{14} \text{Hz}) = 9.95 \times 10^{-20} \frac{\text{J}}{\text{photon}}.$$
21.3

The same process above can be used to determine the energy associated with one photon of representative visible light. According to the first figure in this section, one representative wavelength for visible light is 500 nm.

$$f = \frac{c}{\lambda} = \frac{3.00 \times 10^8 \,\text{m/s}}{5.00 \times 10^{-7} \,\text{m}} = 6.00 \times 10^{14} \,\text{Hz}.$$
21.4

$$\frac{E}{n} = h \cdot f = (6.63 \times 10^{-34} \text{J} \cdot \text{s}) (6.00 \times 10^{14} \text{Hz}) = 3.98 \times 10^{-19} \frac{\text{J}}{\text{photon}}.$$
 21.5

Discussion

This example verifies that as the wavelength of light decreases, the quantum energy increases. This explains why a fire burning with a blue flame is considered more dangerous than a fire with a red flame. Each photon of short-wavelength blue light emitted carries a greater amount of energy than a long-wavelength red light. This example also helps explain the differences in the 3,000 K, 4,000 K, and 6,000 K lines shown in the first figure in this section. As the temperature is increased, more energy is available for a greater number of short-wavelength photons to be emitted.

Practice Problems

- 1. An AM radio station broadcasts at a frequency of 1,530 kHz . What is the energy in Joules of a photon emitted from this station?
 - a. $10.1 \times 10^{-26} \text{ J}$
 - b. 1.01 × 10⁻²⁸ J
 - c. 1.01×10^{-29} J
 - d. 1.01×10^{-27} J
- 2. A photon travels with energy of 1.0 eV. What type of EM radiation is this photon?
 - a. visible radiation

- b. microwave radiation
- c. infrared radiation
- d. ultraviolet radiation

Check Your Understanding

- 3. Do reflective or absorptive surfaces more closely model a perfect blackbody?
 - a. reflective surfaces
 - b. absorptive surfaces
- **4**. A black T-shirt is a good model of a blackbody. However, it is not perfect. What prevents a black T-shirt from being considered a perfect blackbody?
 - a. The T-shirt reflects some light.
 - b. The T-shirt absorbs all incident light.
 - c. The T-shirt re-emits all the incident light.
 - d. The T-shirt does not reflect light.
- 5. What is the mathematical relationship linking the energy of a photon to its frequency?
 - a. $E = \frac{hf}{n}$
 - a. $E = \frac{n}{n}$ b. $E = \frac{nh}{f}$
 - c. $E = \frac{\int_{h}^{h}}{h}$
 - d. E = nhf
- 6. Why do we not notice quantization of photons in everyday experience?
 - a. because the size of each photon is very large
 - b. because the mass of each photon is so small
 - c. because the energy provided by photons is very large
 - d. because the energy provided by photons is very small
- 7. Two flames are observed on a stove. One is red while the other is blue. Which flame is hotter?
 - a. The red flame is hotter because red light has lower frequency.
 - b. The red flame is hotter because red light has higher frequency.
 - c. The blue flame is hotter because blue light has lower frequency.
 - d. The blue flame is hotter because blue light has higher frequency.
- **8**. Your pupils dilate when visible light intensity is reduced. Does wearing sunglasses that lack UV blockers increase or decrease the UV hazard to your eyes? Explain.
 - a. Increase, because more high-energy UV photons can enter the eye.
 - b. Increase, because less high-energy UV photons can enter the eye.
 - c. Decrease, because more high-energy UV photons can enter the eye.
 - d. Decrease, because less high-energy UV photons can enter the eye.
- **9**. The temperature of a blackbody radiator is increased. What will happen to the most intense wavelength of light emitted as this increase occurs?
 - a. The wavelength of the most intense radiation will vary randomly.
 - b. The wavelength of the most intense radiation will increase.
 - c. The wavelength of the most intense radiation will remain unchanged.
 - d. The wavelength of the most intense radiation will decrease.

21.2 Einstein and the Photoelectric Effect

Section Learning Objectives

By the end of this section, you will be able to do the following:

- Describe Einstein's explanation of the photoelectric effect
- Describe how the photoelectric effect could not be explained by classical physics
- Calculate the energy of a photoelectron under given conditions
- Describe use of the photoelectric effect in biological applications, photoelectric devices and movie soundtracks

Section Key Terms

electric eye photoelectric effect photoelectron photon

The Photoelectric Effect

Teacher Support

[EL]Ask the students what they think the term *photoelectric* means. How does the term relate to its definition?

When light strikes certain materials, it can eject electrons from them. This is called the **photoelectric effect**, meaning that light (*photo*) produces electricity. One common use of the photoelectric effect is in light meters, such as those that adjust the automatic iris in various types of cameras. Another use is in solar cells, as you probably have in your calculator or have seen on a rooftop or a roadside sign. These make use of the photoelectric effect to convert light into electricity for running different devices.



Figure 21.5 The photoelectric effect can be observed by allowing light to fall on the metal plate in this evacuated tube. Electrons ejected by the light are collected on the collector wire and measured as a current. A retarding voltage between the collector wire and plate can then be adjusted so as to determine the energy of the ejected electrons. (credit: P. P. Urone)

Revolutionary Properties of the Photoelectric Effect

When Max Planck theorized that energy was quantized in a blackbody radiator, it is unlikely that he would have recognized just how revolutionary his idea was. Using tools similar to the light meter in <u>Figure 21.5</u>, it would take a scientist of Albert Einstein's stature to fully discover the implications of Max Planck's radical concept.

Through careful observations of the photoelectric effect, Albert Einstein realized that there were several characteristics that could be explained only if *EM radiation is itself quantized*. While these characteristics will be explained a bit later in this section, you can already begin to appreciate why Einstein's idea is very important. It means that the apparently continuous stream of energy in an EM wave is actually not a continuous stream at all. In fact, the EM wave itself is actually composed of tiny quantum packets of energy called **photons**.

In equation form, Einstein found the energy of a photon or **photoelectron** to be

E = hf,

where *E* is the energy of a photon of frequency *f* and *h* is Planck's constant. A beam from a flashlight, which to this point had been considered a wave, instead could now be viewed as a series of photons, each providing a specific amount of energy see <u>Figure 21.6</u>. Furthermore, the amount of energy within each individual photon is based upon its individual frequency, as

dictated by E = hf. As a result, the total amount of energy provided by the beam could now be viewed as the sum of all frequency-dependent photon energies added together.



Figure 21.6 An EM wave of frequency *f* is composed of photons, or individual quanta of EM radiation. The energy of each photon is E = hf, where *h* is Planck's constant and *f* is the frequency of the EM radiation. Higher intensity means more photons per unit area per second. The flashlight emits large numbers of photons of many different frequencies, hence others have energy E' = hf', and so on.

Just as with Planck's blackbody radiation, Einstein's concept of the photon could take hold in the scientific community only if it could succeed where classical physics failed. The photoelectric effect would be a key to demonstrating Einstein's brilliance.

Consider the following five properties of the photoelectric effect. All of these properties are consistent with the idea that individual photons of EM radiation are absorbed by individual electrons in a material, with the electron gaining the photon's energy. Some of these properties are inconsistent with the idea that EM radiation is a simple wave. For simplicity, let us consider what happens with monochromatic EM radiation in which all photons have the same energy *hf*.



Figure 21.7 Incident radiation strikes a clean metal surface, ejecting multiple electrons from it. The manner in which the frequency and intensity of the incoming radiation affect the ejected electrons strongly suggests that electromagnetic radiation is quantized. This event, called the photoelectric effect, is strong evidence for the existence of photons.

- 1. If we vary the frequency of the EM radiation falling on a clean metal surface, we find the following: For a given material, there is a threshold frequency f_0 for the EM radiation below which no electrons are ejected, regardless of intensity. Using the photon model, the explanation for this is clear. Individual photons interact with individual electrons. Thus if the energy of an individual photon is too low to break an electron away, no electrons will be ejected. However, if EM radiation were a simple wave, sufficient energy could be obtained simply by increasing the intensity.
- 2. Once EM radiation falls on a material, electrons are ejected without delay. As soon as an individual photon of sufficiently high frequency is absorbed by an individual electron, the electron is ejected. If the EM radiation were a simple wave, several minutes would be required for sufficient energy to be deposited at the metal surface in order to eject an electron.
- 3. The number of electrons ejected per unit time is proportional to the intensity of the EM radiation and to no other characteristic. High-intensity EM radiation consists of large numbers of photons per unit area, with all photons having the same characteristic energy, *hf*. The increased number of photons per unit area results in an increased number of electrons per unit area ejected.
- 4. If we vary the intensity of the EM radiation and measure the energy of ejected electrons, we find the following: *The maximum kinetic energy of ejected electrons is independent of the intensity of the EM radiation*. Instead, as noted in point 3 above, increased intensity results in more electrons of the same energy being ejected. If EM radiation were a simple wave, a higher intensity could transfer more energy, and higher-energy electrons would be ejected.
- 5. The kinetic energy KE of an ejected electron equals the photon energy minus the binding energy BE of the electron in the

specific material. An individual photon can give all of its energy to an electron. The photon's energy is partly used to break the electron away from the material. The remainder goes into the ejected electron's kinetic energy. In equation form, this is given by

where KE_e is the maximum kinetic energy of the ejected electron, hf is the photon's energy, and BE is the binding energy of the electron to the particular material. This equation explains the properties of the photoelectric effect quantitatively and demonstrates that BE is the minimum amount of energy necessary to eject an electron. If the energy supplied is less than BE, the electron cannot be ejected. The binding energy can also be written as $BE = hf_0$, where f_0 is the threshold frequency for the particular material. Figure 21.8 shows a graph of maximum KE_e versus the frequency of incident EM radiation falling on a particular material.



Figure 21.8 A graph of the kinetic energy of an ejected electron, KE_e , versus the frequency of EM radiation impinging on a certain material. There is a threshold frequency below which no electrons are ejected, because the individual photon interacting with an individual electron has insufficient energy to break it away. Above the threshold energy, KE_e increases linearly with *f*, consistent with $KE_e = hf - BE$. The slope of this line is *h*, so the data can be used to determine Planck's constant experimentally.

TIPS FOR SUCCESS

The following five pieces of information can be difficult to follow without some organization. It may be useful to create a table of expected results of each of the five properties, with one column showing the classical wave model result and one column showing the modern photon model result.

The table may look something like Table 21.1

Threshold Frequency	
Electron Ejection Delay	
Intensity of EM Radiation	
Speed of Ejected Electrons	
Relationship between Kinetic Energy and Binding Energy	

Table 21.1 Table of Expected Results

Virtual Physics

Photoelectric Effect

Click to view content (http://www.openstax.org/l/28photoelectric)

In this demonstration, see how light knocks electrons off a metal target, and recreate the experiment that spawned the field of quantum mechanics.

GRASP CHECK

In the circuit provided, what are the three ways to increase the current?

- a. decrease the intensity, decrease the frequency, alter the target
- b. decrease the intensity, decrease the frequency, don't alter the target
- c. increase the intensity, increase the frequency, alter the target
- d. increase the intensity, increase the frequency, alter the target

WORKED EXAMPLE

Photon Energy and the Photoelectric Effect: A Violet Light

(a) What is the energy in joules and electron volts of a photon of 420-nm violet light? (b) What is the maximum kinetic energy of electrons ejected from calcium by 420 nm violet light, given that the binding energy of electrons for calcium metal is 2.71 eV?

Strategy

To solve part (a), note that the energy of a photon is given by E = hf. For part (b), once the energy of the photon is calculated, it is a straightforward application of $KE_e = hf - BE$ to find the ejected electron's maximum kinetic energy, since BE is given.

Solution for (a)

Photon energy is given by

$$E = hf$$
.

Since we are given the wavelength rather than the frequency, we solve the familiar relationship $c = f \lambda$ for the frequency, yielding

$$f = \frac{c}{\lambda}$$
 21.7

Combining these two equations gives the useful relationship

$$E = \frac{hc}{\lambda}.$$
 21.8

Now substituting known values yields

$$E = \frac{(6.63 \times 10^{-34} \text{J} \cdot \text{s})(3.00 \times 10^8 \text{m/s})}{4.20 \times 10^{-7} \text{m}} = 4.74 \times 10^{-19} \text{J}.$$
21.9

Converting to eV, the energy of the photon is

$$E = (4.74 \times 10^{-19} \text{J} \cdot \text{s}) \frac{1 \text{ eV}}{1.60 \times 10^{-19} \text{J}} = 2.96 \text{ eV}.$$
21.10

Solution for (b)

Finding the kinetic energy of the ejected electron is now a simple application of the equation $KE_e = hf - BE$. Substituting the photon energy and binding energy yields

$$KE_e = hf - BE = 2.96 \text{ eV} - 2.71 \text{ eV} = 0.25 \text{ eV}.$$
 21.11

Discussion

The energy of this 420 nm photon of violet light is a tiny fraction of a joule, and so it is no wonder that a single photon would be difficult for us to sense directly—humans are more attuned to energies on the order of joules. But looking at the energy in electron volts, we can see that this photon has enough energy to affect atoms and molecules. A DNA molecule can be broken with about 1 eV of energy, for example, and typical atomic and molecular energies are on the order of eV, so that the photon in this example could have biological effects, such as sunburn. The ejected electron has rather low energy, and it would not travel far,

except in a vacuum. The electron would be stopped by a retarding potential of only 0.26 eV, a slightly larger KE than calculated above. In fact, if the photon wavelength were longer and its energy less than 2.71 eV, then the formula would give a negative kinetic energy, an impossibility. This simply means that the 420 nm photons with their 2.96 eV energy are not much above the frequency threshold. You can see for yourself that the threshold wavelength is 458 nm (blue light). This means that if calcium metal were used in a light meter, the meter would be insensitive to wavelengths longer than those of blue light. Such a light meter would be completely insensitive to red light, for example.

Practice Problems

- **10.** What is the longest-wavelength EM radiation that can eject a photoelectron from silver, given that the bonding energy is 4.73 eV? Is this radiation in the visible range?
 - a. 2.63×10^{-7} m; No, the radiation is in microwave region.
 - b. 2.63×10^{-7} m; No, the radiation is in visible region.
 - c. 2.63×10^{-7} m; No, the radiation is in infrared region.
 - d. 2.63×10^{-7} m; No, the radiation is in ultraviolet region.
- **11**. What is the maximum kinetic energy in eV of electrons ejected from sodium metal by 450-nm EM radiation, given that the binding energy is 2.28 eV?
 - a. 0.48 V
 - b. 0.82 eV
 - c. 1.21 eV
 - d. 0.48 eV

Technological Applications of the Photoelectric Effect

While Einstein's understanding of the photoelectric effect was a transformative discovery in the early 1900s, its presence is ubiquitous today. If you have watched streetlights turn on automatically in response to the setting sun, stopped elevator doors from closing simply by putting your hands between them, or turned on a water faucet by sliding your hands near it, you are familiar with the **electric eye**, a name given to a group of devices that use the photoelectric effect for detection.

All these devices rely on photoconductive cells. These cells are activated when light is absorbed by a semi-conductive material, knocking off a free electron. When this happens, an electron void is left behind, which attracts a nearby electron. The movement of this electron, and the resultant chain of electron movements, produces a current. If electron ejection continues, further holes are created, thereby increasing the electrical conductivity of the cell. This current can turn switches on and off and activate various familiar mechanisms.

One such mechanism takes place where you may not expect it. Next time you are at the movie theater, pay close attention to the sound coming out of the speakers. This sound is actually created using the photoelectric effect! The audiotape in the projector booth is a transparent piece of film of varying width. This film is fed between a photocell and a bright light produced by an exciter lamp. As the transparent portion of the film varies in width, the amount of light that strikes the photocell varies as well. As a result, the current in the photoconductive circuit changes with the width of the filmstrip. This changing current is converted to a changing frequency, which creates the soundtrack commonly heard in the theater.

WORK IN PHYSICS

Solar Energy Physicist

According to the U.S. Department of Energy, Earth receives enough sunlight each hour to power the entire globe for a year. While converting all of this energy is impossible, the job of the solar energy physicist is to explore and improve upon solar energy conversion technologies so that we may harness more of this abundant resource.

The field of solar energy is not a new one. For over half a century, satellites and spacecraft have utilized photovoltaic cells to create current and power their operations. As time has gone on, scientists have worked to adapt this process so that it may be used in homes, businesses, and full-scale power stations using solar cells like the one shown in Figure 21.9.



Figure 21.9 A solar cell is an example of a photovoltaic cell. As light strikes the cell, the cell absorbs the energy of the photons. If this energy exceeds the binding energy of the electrons, then electrons will be forced to move in the cell, thereby producing a current. This current may be used for a variety of purposes. (credit: U.S. Department of Energy)

Solar energy is converted to electrical energy in one of two manners: direct transfer through photovoltaic cells or thermal conversion through the use of a CSP, concentrating solar power, system. Unlike electric eyes, which trip a mechanism when current is lost, photovoltaic cells utilize semiconductors to directly transfer the electrons released through the photoelectric effect into a directed current. The energy from this current can then be converted for storage, or immediately used in an electric process. A CSP system is an indirect method of energy conversion. In this process, light from the Sun is channeled using parabolic mirrors. The light from these mirrors strikes a thermally conductive material, which then heats a pool of water. This water, in turn, is converted to steam, which turns a turbine and creates electricity. While indirect, this method has long been the traditional means of large-scale power generation.

There are, of course, limitations to the efficacy of solar power. Cloud cover, nightfall, and incident angle strike at high altitudes are all factors that directly influence the amount of light energy available. Additionally, the creation of photovoltaic cells requires rare-earth minerals that can be difficult to obtain. However, the major role of a solar energy physicist is to find ways to improve the efficiency of the solar energy conversion process. Currently, this is done by experimenting with new semi conductive materials, by refining current energy transfer methods, and by determining new ways of incorporating solar structures into the current power grid.

Additionally, many solar physicists are looking into ways to allow for increased solar use in impoverished, more remote locations. Because solar energy conversion does not require a connection to a large-scale power grid, research into thinner, more mobile materials will permit remote cultures to use solar cells to convert sunlight collected during the day into stored energy that can then be used at night.

Regardless of the application, solar energy physicists are an important part of the future in responsible energy growth. While a doctoral degree is often necessary for advanced research applications, a bachelor's or master's degree in a related science or engineering field is typically enough to gain access into the industry. Computer skills are very important for energy modeling, including knowledge of CAD software for design purposes. In addition, the ability to collaborate and communicate with others is critical to becoming a solar energy physicist.

GRASP CHECK

What role does the photoelectric effect play in the research of a solar energy physicist?

- a. The understanding of photoelectric effect allows the physicist to understand the generation of light energy when using photovoltaic cells.
- b. The understanding of photoelectric effect allows the physicist to understand the generation of electrical energy when using photovoltaic cells.
- c. The understanding of photoelectric effect allows the physicist to understand the generation of electromagnetic energy when using photovoltaic cells.
- d. The understanding of photoelectric effect allows the physicist to understand the generation of magnetic energy when using photovoltaic cells.

Check Your Understanding

- 12. How did Einstein's model of photons change the view of a beam of energy leaving a flashlight?
 - a. A beam of light energy is now considered a continual stream of wave energy, not photons.
 - b. A beam of light energy is now considered a collection of photons, each carrying its own individual energy.
- 13. True or false—Visible light is the only type of electromagnetic radiation that can cause the photoelectric effect.
 - a. false
 - b. true
- 14. Is the photoelectric effect a direct consequence of the wave character of EM radiation or the particle character of EM radiation?
 - a. The photoelectric effect is a direct consequence of the particle nature of EM radiation.
 - b. The photoelectric effect is a direct consequence of the wave nature of EM radiation.
 - c. The photoelectric effect is a direct consequence of both the wave and particle nature of EM radiation.
 - d. The photoelectric effect is a direct consequence of neither the wave nor the particle nature of EM radiation.
- 15. Which aspects of the photoelectric effect can only be explained using photons?
 - a. aspects 1, 2, and 3
 - b. aspects 1, 2, and 4
 - c. aspects 1, 2, 4 and 5
 - d. aspects 1, 2, 3, 4 and 5
- 16. In a photovoltaic cell, what energy transformation takes place?
 - a. Solar energy transforms into electric energy.
 - b. Solar energy transforms into mechanical energy.
 - c. Solar energy transforms into thermal energy.
 - d. In a photovoltaic cell, thermal energy transforms into electric energy.
- 17. True or false—A current is created in a photoconductive cell, even if only one electron is expelled from a photon strike.
 - a. false
 - b. true
- 18. What is a photon and how is it different from other fundamental particles?
 - a. A photon is a quantum packet of energy; it has infinite mass.
 - b. A photon is a quantum packet of energy; it is massless.
 - c. A photon is a fundamental particle of an atom; it has infinite mass.
 - d. A photon is a fundamental particle of an atom; it is massless.

21.3 The Dual Nature of Light

Section Learning Objectives

By the end of this section, you will be able to do the following:

- Describe the Compton effect
- Calculate the momentum of a photon
- Explain how photon momentum is used in solar sails
- Explain the particle-wave duality of light

Section Key Terms

Compton effect particle-wave duality photon momentum

Photon Momentum

Do photons abide by the fundamental properties of physics? Can packets of electromagnetic energy possibly follow the same rules as a ping-pong ball or an electron? Although strange to consider, the answer to both questions is yes.

Despite the odd nature of photons, scientists prior to Einstein had long suspected that the fundamental particle of

electromagnetic radiation shared properties with our more macroscopic particles. This is no clearer than when considering the photoelectric effect, where photons knock electrons out of a substance. While it is strange to think of a massless particle exhibiting momentum, it is now a well-established fact within the scientific community. Figure 21.10 shows macroscopic evidence of **photon momentum**.



Figure 21.10 The tails of the Hale-Bopp comet point away from the Sun, evidence that light has momentum. Dust emanating from the body of the comet forms this tail. Particles of dust are pushed away from the Sun by light reflecting from them. The blue, ionized gas tail is also produced by photons interacting with atoms in the comet material. (credit: Geoff Chester, U.S. Navy, via Wikimedia Commons)

<u>Figure 21.10</u> shows a comet with two prominent tails. Comet tails are composed of gases and dust evaporated from the body of the comet and ionized gas. What most people do not know about the tails is that they always point *away* from the Sun rather than trailing behind the comet. This can be seen in the diagram.

Why would this be the case? The evidence indicates that the dust particles of the comet are forced away from the Sun when photons strike them. Evidently, photons carry momentum in the direction of their motion away from the Sun, and some of this momentum is transferred to dust particles in collisions. The blue tail is caused by the solar wind, a stream of plasma consisting primarily of protons and electrons evaporating from the corona of the Sun.

Momentum, The Compton Effect, and Solar Sails

Momentum is conserved in quantum mechanics, just as it is in relativity and classical physics. Some of the earliest direct experimental evidence of this came from the scattering of X-ray photons by electrons in substances, a phenomenon discovered by American physicist Arthur H. Compton (1892–1962). Around 1923, Compton observed that X-rays reflecting from materials had decreased energy and correctly interpreted this as being due to the scattering of the X-ray photons by electrons. This phenomenon could be handled as a collision between two particles—a photon and an electron at rest in the material. After careful observation, it was found that both energy and momentum were conserved in the collision. See Figure 21.11. For the discovery of this conserved scattering, now known as the **Compton effect**, Arthur Compton was awarded the Nobel Prize in 1929.

Shortly after the discovery of Compton scattering, the value of the photon momentum, $\mathbf{p} = \frac{h}{2}$,

was determined by Louis de Broglie. In this equation, called the de Broglie relation, h represents Planck's constant and λ is the photon wavelength.



Figure 21.11 The Compton effect is the name given to the scattering of a photon by an electron. Energy and momentum are conserved, resulting in a reduction of both for the scattered photon.

We can see that photon momentum is small, since $\mathbf{p} = h/\lambda$. and *h* is very small. It is for this reason that we do not ordinarily observe photon momentum. Our mirrors do not recoil when light reflects from them, except perhaps in cartoons. Compton saw the effects of photon momentum because he was observing X-rays, which have a small wavelength and a relatively large momentum, interacting with the lightest of particles, the electron.

WORKED EXAMPLE

Electron and Photon Momentum Compared

(a) Calculate the momentum of a visible photon that has a wavelength of 500 nm. (b) Find the velocity of an electron having the same momentum. (c) What is the energy of the electron, and how does it compare with the energy of the photon?

Strategy

Finding the photon momentum is a straightforward application of its definition: $\mathbf{p} = h/\lambda$. If we find the photon momentum is small, we can assume that an electron with the same momentum will be nonrelativistic, making it easy to find its velocity and kinetic energy from the classical formulas.

Solution for (a)

Photon momentum is given by the de Broglie relation.

$$\mathbf{p} = \frac{h}{\lambda}$$
 21.12

Entering the given photon wavelength yields

$$\mathbf{p} = \frac{6.63 \times 10^{-34} J \cdot s}{5.00 \times 10^{-7} \mathrm{m}} = 1.33 \times 10^{-27} \mathrm{kg} \cdot \mathrm{m/s}.$$
 21.13

Solution for (b)

Since this momentum is indeed small, we will use the classical expression p = mv to find the velocity of an electron with this momentum. Solving for *v* and using the known value for the mass of an electron gives

$$v = \frac{p}{m} = \frac{1.33 \times 10^{-27} \text{kg} \cdot \text{m/s}}{9.11 \times 10^{-31} \text{kg}} = 1,459.9 \text{ m/s} \approx 1,460 \text{ m/s}.$$
21.14

Solution for (c)

The electron has kinetic energy, which is classically given by

$$KE_e = \frac{1}{2}mv^2.$$

Thus,

$$KE_e = \frac{1}{2}(9.11 \times 10^{-31} \text{kg})(1,456 \text{m/s})^2 = 9.64 \times 10^{-25} \text{J}.$$
 21.16

Converting this to eV by multiplying by $\frac{(1 \text{ eV})}{(1.602 \times 10^{-19} \text{ J})}$ yields

$$KE_e = 6.02 \times 10^{-6} \text{eV}.$$
 21.17

The photon energy *E* is

$$E = \frac{hc}{\lambda} = \frac{(6.63 \times 10^{-34} \text{ J} \cdot \text{s}) (3.00 \times 10^8 \text{ m/s})}{5.00 \times 10^{-7} \text{ m}} = 3.98 \times 10^{-19} \text{ J} = 2.48 \text{ eV},$$
21.18

which is about five orders of magnitude greater.

Discussion

Even in huge numbers, the total momentum that photons carry is small. An electron that carries the same momentum as a 500-nm photon will have a 1,460 m/s velocity, which is clearly nonrelativistic. This is borne out by the experimental observation that it takes far less energy to give an electron the same momentum as a photon. That said, for high-energy photons interacting with small masses, photon momentum may be significant. Even on a large scale, photon momentum can have an effect if there

are enough of them and if there is nothing to prevent the slow recoil of matter. Comet tails are one example, but there are also proposals to build space sails that use huge low-mass mirrors (made of aluminized Mylar) to reflect sunlight. In the vacuum of space, the mirrors would gradually recoil and could actually accelerate spacecraft within the solar system. See the following figure.

TIPS FOR SUCCESS

When determining energies in particle physics, it is more sensible to use the unit eV instead of Joules. Using eV will help you to recognize differences in magnitude more easily and will make calculations simpler. Also, eV is used by scientists to describe the binding energy of particles and their rest mass, so using eV will eliminate the need to convert energy quantities. Finally, eV is a convenient unit when linking electromagnetic forces to particle physics, as one eV is the amount energy given to an electron when placed in a field of 1-V potential difference.

Practice Problems

19. Find the momentum of a 4.00-cm wavelength microwave photon.

- a. $0.83 \times 10^{-32} \text{ kg} \cdot \text{m/s}$
- b. $1.66 \times 10^{-34} \text{ kg} \cdot \text{m/s}$
- c. $0.83 \times 10^{-34} \text{ kg} \cdot \text{m/s}$
- d. $1.66 \times 10^{-32} \text{ kg} \cdot \text{m/s}$

20. Calculate the wavelength of a photon that has the same momentum of a proton moving at 1.00 percent of the speed of light.

- a. 2.43×10^{-10} m
- b. $2.43 \times 10^{-12} \text{ m}$
- c. 1.32×10^{-15} m
- d. $1.32 \times 10^{-13} \text{ m}$



Figure 21.12 (a) Space sails have been proposed that use the momentum of sunlight reflecting from gigantic low-mass sails to propel spacecraft about the solar system. A Russian test model of this (the Cosmos 1) was launched in 2005, but did not make it into orbit due to a rocket failure. (b) A U.S. version of this, labeled LightSail-1, is scheduled for trial launches in 2016. It will have a 40 m2 sail. (credit: Kim Newton/NASA)

LightSail-1 Project

"Provide ships or sails adapted to the heavenly breezes, and there will be some who will brave even that void."

— Johannes Kepler (in a letter to Galileo Galilei in 1608)



Figure 21.13 NASA's NanoSail-D, a precursor to LightSail-1, with its sails deployed. The Planetary Society will be launching LightSail-1 in early 2016. (credit: NASA/MSFC/D, Wikimedia Commons)

Traversing the Solar System using nothing but the Sun's power has long been a fantasy of scientists and science fiction writers alike. Though physicists like Compton, Einstein, and Planck all provided evidence of light's propulsive capacity, it is only recently that the technology has become available to truly put these visions into motion. In 2016, by sending a lightweight satellite into space, the LightSail-1 project is designed to do just that.

A citizen-funded project headed by the Planetary Society, the 5.45-million-dollar LightSail-1 project is set to launch two crafts into orbit around the Earth. Each craft is equipped with a 32-square-meter solar sail prepared to unfurl once a rocket has launched it to an appropriate altitude. The sails are made of large mirrors, each a quarter of the thickness of a trash bag, which will receive an impulse from the Sun's reflecting photons. Each time the Sun's photon strikes the craft's reflective surface and bounces off, it will provide a momentum to the sail much greater than if the photon were simply absorbed.

Attached to three tiny satellites called CubeSats, whose combined volume is no larger than a loaf of bread, the received momentum from the Sun's photons should be enough to record a substantial increase in orbital speed. The intent of the LightSail-1 mission is to prove that the technology behind photon momentum travel is sound and can be done cheaply. A test flight in May 2015 showed that the craft's Mylar sails could unfurl on command. With another successful result in 2016, the Planetary Society will be planning future versions of the craft with the hopes of eventually achieving interplanetary satellite travel. Though a few centuries premature, Kepler's fantastic vision may not be that far away.

If eventually set into interplanetary launch, what will be the effect of continual photon bombardment on the motion of a craft similar to LightSail-1?

- a. It will result in continual acceleration of the craft.
- b. It will first accelerate and then decelerate the craft.
- c. It will first decelerate and then accelerate the craft.
- d. It will result in the craft moving at constant velocity.

Particle-Wave Duality

We have long known that EM radiation is like a wave, capable of interference and diffraction. We now see that light can also be modeled as particles—massless photons of discrete energy and momentum. We call this twofold nature the **particle-wave duality**, meaning that EM radiation has properties of both particles and waves. This may seem contradictory, since we ordinarily deal with large objects that never act like both waves and particles. An ocean wave, for example, looks nothing like a grain of sand. However, this so-called duality is simply a term for properties of the photon analogous to phenomena we can observe directly, on a macroscopic scale. See <u>Figure 21.14</u>. If this term seems strange, it is because we do not ordinarily observe details on the quantum level directly, and our observations yield either particle-like *or* wave-like properties, but never both simultaneously.



Figure 21.14 (a) The interference pattern for light through a double slit is a wave property understood by analogy to water waves. (b) The properties of photons having quantized energy and momentum and acting as a concentrated unit are understood by analogy to macroscopic particles.

Since we have a particle-wave duality for photons, and since we have seen connections between photons and matter in that both have momentum, it is reasonable to ask whether there is a particle-wave duality for matter as well. If the EM radiation we once thought to be a pure wave has particle properties, is it possible that matter has wave properties? The answer, strangely, is yes. The consequences of this are tremendous, as particle-wave duality has been a constant source of scientific wonder during the twentieth and twenty-first centuries.

Check Your Understanding

- 21. What fundamental physics properties were found to be conserved in Compton scattering?
 - a. energy and wavelength
 - b. energy and momentum
 - c. mass and energy
 - d. energy and angle
- **22.** Why do classical or relativistic momentum equations not work in explaining the conservation of momentum that occurs in Compton scattering?
 - a. because neither classical nor relativistic momentum equations utilize mass as a variable in their equations
 - b. because relativistic momentum equations utilize mass as a variable in their formulas but classical momentum equations do not
 - c. because classical momentum equations utilize mass as a variable in their formulas but relativistic momentum equations do not
 - d. because both classical and relativistic momentum equations utilize mass as a variable in their formulas
- 23. If solar sails were constructed with more massive materials, how would this influence their effectiveness?
 - a. The effect of the momentum would increase due to the decreased inertia of the sails.
 - b. The effect of the momentum would reduce due to the decreased inertia of the sails.
 - c. The effect of the momentum would increase due to the increased inertia of the sails.
 - d. The effect of the momentum would be reduced due to the increased inertia of the sails.
- 24. True or false—It is possible to propel a solar sail craft using just particles within the solar wind.
 - a. true
 - b. false
- 25. True or false—Photon momentum more directly supports the wave model of light.
 - a. false
 - b. true

- **26**. True or false—wave-particle duality exists for objects on the macroscopic scale.
 - a. false
 - b. true
- 27. What type of electromagnetic radiation was used in Compton scattering?
 - a. visible light
 - b. ultraviolet radiation
 - c. radio waves
 - d. X-rays

KEY TERMS

- **blackbody** object that absorbs all radiated energy that strikes it and also emits energy across all wavelengths of the electromagnetic spectrum
- **Compton effect** phenomenon whereby X-rays scattered from materials have decreased energy
- **electric eye** group of devices that use the photoelectric effect for detection
- **particle-wave duality** property of behaving like either a particle or a wave; the term for the phenomenon that all particles have wave-like characteristics and waves have particle-like characteristics
- **photoelectric effect** phenomenon whereby some materials eject electrons when exposed to light

photoelectron electron that has been ejected from a

SECTION SUMMARY

21.1 Planck and Quantum Nature of Light

- A blackbody will radiate energy across all wavelengths of the electromagnetic spectrum.
- Radiation of a blackbody will peak at a particular wavelength, dependent on the temperature of the blackbody.
- Analysis of blackbody radiation led to the field of quantum mechanics, which states that radiated energy can only exist in discrete quantum states.

21.2 Einstein and the Photoelectric Effect

- The photoelectric effect is the process in which EM radiation ejects electrons from a material.
- Einstein proposed photons to be quanta of EM radiation having energy E = hf, where *f* is the frequency of the radiation.
- All EM radiation is composed of photons. As Einstein

KEY EQUATIONS

21.1 Planck and Quantum Nature of Light

quantum energy E = nhf

21.2 Einstein and the Photoelectric Effect

energy of a photon

$$E = hf$$

material by a photon of light

- **photon** a quantum, or particle, of electromagnetic radiation
- **photon momentum** amount of momentum of a photon, calculated by $\mathbf{p} = \frac{h}{\lambda}$
- **quantized** the fact that certain physical entities exist only with particular discrete values and not every conceivable value
- **quantum** discrete packet or bundle of a physical entity such as energy
- **ultraviolet catastrophe** misconception that blackbodies would radiate high frequency energy at a much higher rate than energy radiated at lower frequencies

explained, all characteristics of the photoelectric effect are due to the interaction of individual photons with individual electrons.

• The maximum kinetic energy KE_e of ejected electrons (photoelectrons) is given by $KE_e = hf - BE$, where hfis the photon energy and BE is the binding energy (or work function) of the electron in the particular material.

21.3 The Dual Nature of Light

- Compton scattering provided evidence that photonelectron interactions abide by the principles of conservation of momentum and conservation of energy.
- The momentum of individual photons, quantified by $\mathbf{p} = \frac{h}{\lambda}$, can be used to explain observations of comets and may lead to future space technologies.
- Electromagnetic waves and matter have both wave-like and particle-like properties. This phenomenon is defined as particle-wave duality.

maximum kinetic energy of a photoelectron	$KE_e = hf - BE$

binding energy of an electron $BE = hf_O$

21.3 The Dual Nature of Light

momentum of a photon (deBroglie relation)

$$\mathbf{p} = \frac{h}{\lambda}$$

CHAPTER REVIEW Concept Items

21.1 Planck and Quantum Nature of Light

- 1. What aspect of the blackbody spectrum forced Planck to propose quantization of energy levels in atoms and molecules?
 - a. Radiation occurs at a particular frequency that does not change with the energy supplied.
 - b. Certain radiation occurs at a particular frequency that changes with the energy supplied.
 - c. Maximum radiation would occur at a particular frequency that does not change with the energy supplied.
 - d. Maximum radiation would occur at a particular frequency that changes with the energy supplied.
- **2**. Two lasers shine red light at 650 nm. One laser is twice as bright as the other. Explain this difference using photons and photon energy.
 - a. The brighter laser emits twice the number of photons and more energy per photon.
 - b. The brighter laser emits twice the number of photons and less energy per photon.
 - c. Both lasers emit equal numbers of photons and equivalent amounts of energy per photon.
 - d. The brighter laser emits twice the number of photons but both lasers emit equivalent amounts of energy per photon.
- **3.** Consider four stars in the night sky: red, yellow, orange, and blue. The photons of which star will carry the greatest amount of energy?
 - a. blue
 - b. orange
 - c. red
 - d. yellow
- **4**. A lightbulb is wired to a variable resistor. What will happen to the color spectrum emitted by the bulb as the resistance of the circuit is increased?
 - a. The bulb will emit greener light.
 - b. The bulb will emit bluer light.
 - c. The bulb will emit more ultraviolet light.
 - d. The bulb will emit redder light.

21.2 Einstein and the Photoelectric Effect

- 5. Light is projected onto a semi-conductive surface. However, no electrons are ejected. What will happen when the light intensity is increased?
 - a. An increase in light intensity decreases the number of photons. However, no electrons are ejected.

- b. Increase in light intensity increases the number of photons, so electrons with higher kinetic energy are ejected.
- c. An increase in light intensity increases the number of photons, so electrons will be ejected.
- d. An increase in light intensity increases the number of photons. However, no electrons are ejected.
- **6**. True or false—The concept of a work function (or binding energy) is permissible under the classical wave model.
 - a. false
 - b. true
- 7. Can a single microwave photon cause cell damage?
 - a. No, there is not enough energy associated with a single microwave photon to result in cell damage.
 - b. No, there is zero energy associated with a single microwave photon, so it does not result in cell damage.
 - c. Yes, a single microwave photon causes cell damage because it does not have high energy.
 - d. Yes, a single microwave photon causes cell damage because it has enough energy.

21.3 The Dual Nature of Light

- **8**. Why don't we feel the momentum of sunlight when we are on the beach?
 - a. The momentum of a singular photon is incredibly small.
 - b. The momentum is not felt because very few photons strike us at any time, and not all have momentum.
 - c. The momentum of a singular photon is large, but very few photons strike us at any time.
 - d. A large number of photons strike us at any time, and so their combined momentum is incredibly large.
- **9**. If a beam of helium atoms is projected through two slits and onto a screen, will an interference pattern emerge?
 - a. No, an interference pattern will not emerge because helium atoms will strike a variety of locations on the screen.
 - b. No, an interference pattern will not emerge because helium atoms will strike at certain locations on the screen.
 - c. Yes, an interference pattern will emerge because helium atoms will strike a variety of locations on the screen.
 - d. Yes, an interference pattern will emerge because helium atoms will strike at certain locations on the screen.

Critical Thinking Items

21.1 Planck and Quantum Nature of Light

- **10**. Explain why the frequency of a blackbody does not double when the temperature is doubled.
 - a. Frequency is inversely proportional to temperature.
 - b. Frequency is directly proportional to temperature.
 - c. Frequency is directly proportional to the square of temperature.
 - d. Frequency is directly proportional to the fourth power of temperature.
- **11**. Why does the intensity shown in the blackbody radiation graph decrease after its peak frequency is achieved?



- a. Because after reaching the peak frequency, the photons created at a particular frequency are too many for energy intensity to continue to decrease.
- b. Because after reaching the peak frequency, the photons created at a particular frequency are too few for energy intensity to continue to decrease.
- c. Because after reaching the peak frequency, the photons created at a particular frequency are too many for energy intensity to continue to increase.
- d. Because after reaching the peak frequency, the photons created at a particular frequency are too few for energy intensity to continue to increase.
- 12. Shortly after the introduction of photography, it was found that photographic emulsions were more sensitive to blue and violet light than they were to red light. Explain why this was the case.
 - a. Blue-violet light contains greater amount of energy than red light.
 - b. Blue-violet light contains lower amount of energy than red light.
 - c. Both blue-violet light and red light have the same frequency but contain different amounts of energy.
 - d. Blue-violet light frequency is lower than the frequency of red light.

- **13**. Why is it assumed that a perfect absorber of light (like a blackbody) must also be a perfect emitter of light?
 - a. To achieve electrostatic equilibrium with its surroundings
 - b. To achieve thermal equilibrium with its surroundings
 - c. To achieve mechanical equilibrium with its surroundings
 - d. To achieve chemical equilibrium with its surroundings

21.2 Einstein and the Photoelectric Effect

- **14**. Light is projected onto a semi-conductive surface. If the intensity is held constant but the frequency of light is increased, what will happen?
 - a. As frequency is increased, electrons will stop being ejected from the surface.
 - b. As frequency is increased, electrons will begin to be ejected from the surface.
 - c. As frequency is increased, it will have no effect on the electrons being ejected as the intensity is the same.
 - d. As frequency is increased, the rate at which the electrons are being ejected will increase.
- **15**. Why is it important to consider what material to use when designing a light meter? Consider the worked example from Section 21-2 for assistance.
 - a. A light meter should contain material that responds only to high frequency light.
 - b. A light meter should contain material that responds to low frequency light.
 - c. A light meter should contain material that has high binding energy.
 - d. A light meter should contain a material that does not show any photoelectric effect.
- 16. Why does overexposure to UV light often result in sunburn when overexposure to visible light does not? This is why you can get burnt even on a cloudy day.
 - a. UV light carries less energy than visible light and can penetrate our body.
 - b. UV light carries more energy than visible light, so it cannot break bonds at the cellular level.
 - c. UV light carries more energy than visible light and can break bonds at the cellular level.
 - d. UV light carries less energy than visible light and cannot penetrate the human body.
- 17. If you pick up and shake a piece of metal that has electrons in it free to move as a current, no electrons fall out. Yet if you heat the metal, electrons can be boiled off. Explain both of these facts as they relate to the amount and distribution of energy involved with shaking the

object as compared with heating it.

- a. Thermal energy is added to the metal at a much higher rate than energy added due to shaking.
- b. Thermal energy is added to the metal at a much lower rate than energy added due to shaking.
- c. If the thermal energy added is below the binding energy of the electrons, they may be *boiled off*.
- d. If the mechanical energy added is below the binding energy of the electrons, they may be *boiled* off.

21.3 The Dual Nature of Light

- 18. In many macroscopic collisions, a significant amount of kinetic energy is converted to thermal energy. Explain why this is not a concern for Compton scattering.
 - a. Because, photons and electrons do not exist on the molecular level, all energy of motion is considered kinetic energy.
 - b. Because, photons exist on the molecular level while electrons do not exist on the molecular level, all energy of motion is considered kinetic energy.
 - c. Because, electrons exist on the molecular level while photons do not exist on the molecular level, all energy of motion is considered kinetic energy.
 - d. Because, photons and electrons exist on the molecular level, all energy of motion is considered kinetic energy.

Problems

21.1 Planck and Quantum Nature of Light

- **22**. How many X-ray photons per second are created by an X-ray tube that produces a flux of X-rays having a power of 1.00 W? Assume the average energy per photon is 75.0 keV.
 - a. 8.33×10^{15} photons
 - b. 9.1×10^7 photons
 - c. 9.1×10^8 photons
 - d. 8.33×10^{13} photons
- **23**. What is the frequency of a photon produced in a CRT using a 25.0-kV accelerating potential? This is similar to the layout as in older color television sets.
 - a. $6.04 \times 10^{-48} \text{ Hz}$
 - b. $2.77 \times 10^{-48} \text{ Hz}$
 - c. $3.02 \times 10^{18} \text{ Hz}$
 - d. 6.04×10^{18} Hz

21.2 Einstein and the Photoelectric Effect

24. What is the binding energy in eV of electrons in magnesium, if the longest-wavelength photon that can eject electrons is 337 nm?

- **19**. In what region of the electromagnetic spectrum will photons be most effective in accelerating a solar sail?
 - a. ultraviolet rays
 - b. infrared rays
 - c. X-rays
 - d. gamma rays
- **20**. True or false—Electron microscopes can resolve images that are smaller than the images resolved by light microscopes.
 - a. false
 - b. true
- **21.** How would observations of Compton scattering change if ultraviolet light were used in place of X-rays?
 - a. Ultraviolet light carries less energy than X-rays. As a result, Compton scattering would be easier to detect.
 - b. Ultraviolet light carries less energy than X-rays. As a result, Compton scattering would be more difficult to detect.
 - c. Ultraviolet light carries more energy than X-rays. As a result, Compton scattering would be easier to detect.
 - d. Ultraviolet light has higher energy than X-rays. As a result, Compton scattering would be more difficult to detect.
 - a. 7.44×10^{-19} J
 - b. 7.44 × 10⁻⁴⁹ J
 - c. 5.90×10^{-17} J
 - d. $5.90 \times 10^{-19} \text{ J}$
- **25**. Photoelectrons from a material with a binding energy of 2.71 eV are ejected by 420-nm photons. Once ejected, how long does it take these electrons to travel 2.50 cm to a detection device?
 - a. 8.5×10^{-6} s
 - b. $3.5 \times 10^{-7} \, \text{s}$
 - c. 43.5×10^{-9} s
 - d. 8.5×10^{-8} s

21.3 The Dual Nature of Light

- **26.** What is the momentum of a 0.0100-nm-wavelength photon that could detect details of an atom?
 - a. $6.626 \times 10^{-27} \text{ kg} \cdot \text{m/s}$
 - b. $6.626 \times 10^{-32} \text{ kg} \cdot \text{m/s}$
 - c. $6.626 \times 10^{-34} \text{ kg} \cdot \text{m/s}$
 - d. $6.626 \times 10^{-23} \text{ kg} \cdot \text{m/s}$
- **27**. The momentum of light is exactly reversed when reflected straight back from a mirror, assuming negligible recoil of the mirror. Thus the change in

momentum is twice the initial photon momentum. Suppose light of intensity 1.00 kW/m2 reflectsfrom a mirror of area 2.00 m2 each second. Using the most general form of Newton's second law, what is the force on the mirror?

Performance Task

21.3 The Dual Nature of Light

- **28**. Our scientific understanding of light has changed over time. There is evidence to support the wave model of light, just as there is evidence to support the particle model of light.
 - Construct a demonstration that supports the wave model of light. Note—One possible method is to use a piece of aluminum foil, razor blade, and laser to demonstrate wave interference. Can you arrange these materials to create an effective demonstration? In writing, explain how evidence

TEST PREP Multiple Choice

21.1 Planck and Quantum Nature of Light

- **29.** A perfect blackbody is a perfect absorber of energy transferred by what method?
 - a. conduction
 - b. convection
 - c. induction
 - d. radiation
- **30.** Which of the following is a physical entity that is quantized?
 - a. electric charge of an ion
 - b. frequency of a sound
 - c. speed of a car
- **31**. Find the energy in joules of photons of radio waves that leave an FM station that has a 90.0-MHz broadcast frequency.
 - a. 1.8×10^{-25} J
 - b. 1.11 × 10⁻²⁵ J
 - c. 7.1 × 10⁻⁴³ J
 - d. 5.96×10^{-26} J
- **32.** Which region of the electromagnetic spectrum will provide photons of the least energy?
 - a. infrared light
 - b. radio waves
 - c. ultraviolet light
 - d. X-rays
- **33.** A hot, black coffee mug is sitting on a kitchen table in a dark room. Because it cannot be seen, one assumes that

- a. 1.33×10^{-5} N
- b. 1.33 × 10⁻⁶ N
- c. 1.33×10^{-7} N
- d. 1.33×10^{-8} N

from your demonstration supports the wave model of light.

2. Construct a demonstration that supports the particle model of light. Note—One possible method is to use a negatively charged electroscope, zinc plate, and three light sources of different frequencies. A red laser, a desk lamp, and ultraviolet lamp are typically used. Can you arrange these materials to demonstrate the photoelectric effect? In writing, explain how evidence from your demonstration supports the particle model of light.

it is not emitting energy in the form of light. Explain the fallacy in this logic.

- a. Not all heat is in the form of light energy.
- b. Not all light energy falls in the visible portion of the electromagnetic spectrum.
- c. All heat is in the form of light energy.
- d. All light energy falls in the visible portion of the electromagnetic spectrum.
- 34. Given two stars of equivalent size, which will have a greater temperature: a red dwarf or a yellow dwarf? Explain. Note—Our sun is considered a yellow dwarf.
 - a. a yellow dwarf, because yellow light has lower frequency
 - b. a red dwarf, because red light has lower frequency
 - c. a red dwarf, because red light has higher frequencyd. a yellow dwarf, because yellow light has higher
 - frequency

21.2 Einstein and the Photoelectric Effect

- 35. What is a quantum of light called?
 - a. electron
 - b. neutron
 - c. photon
 - d. proton
- **36**. Which of the following observations from the photoelectric effect is not a violation of classical physics?
 - a. Electrons are ejected immediately after impact from light.
 - b. Light can eject electrons from a semi-conductive

material.

- c. Light intensity does not influence the kinetic energy of ejected electrons.
- d. No electrons are emitted if the light frequency is too low.
- **37.** If 5 eV of energy is supplied to an electron with a binding energy of 2.3 eV, with what kinetic energy will the electron be launched?
 - a. 2.3 eV
 - b. 7.3 eV
 - c. 11.5 eV
 - d. 2.7 eV
- **38**. Which of the following terms translates to *light-producing voltage*?
 - a. photoelectric
 - b. quantum mechanics
 - c. photoconductive
 - d. photovoltaic
- **39**. Why is high frequency EM radiation considered more dangerous than long wavelength EM radiation?
 - a. Long wavelength EM radiation photons carry less energy and therefore have greater ability to disrupt materials through the photoelectric effect.
 - b. Long wavelength EM radiation photons carry more energy and therefore have greater ability to disrupt materials through the photoelectric effect.
 - c. High frequency EM radiation photons carry less energy and therefore have lower ability to disrupt materials through the photoelectric effect.
 - d. High frequency EM radiation photons carry more energy and therefore have greater ability to disrupt materials through the photoelectric effect.
- **40**. Why are UV, X-rays, and gamma rays considered ionizing radiation?
 - a. UV, X-rays, and gamma rays are capable of ejecting photons from a surface.
 - b. UV, X-rays, and gamma rays are capable of ejecting neutrons from a surface.
 - c. UV, X-rays, and gamma rays are capable of ejecting

Short Answer

21.1 Planck and Quantum Nature of Light

- **46**. Scientists once assumed that all frequencies of light were emitted with equal probability. Explain what the blackbody radiation curve would look like if this were the case.
 - a. The blackbody radiation curve would look like a circular path.
 - b. The blackbody radiation curve would look like an

protons from a surface.

d. UV, X-rays, and gamma rays are capable of ejecting electrons from a surface.

21.3 The Dual Nature of Light

- 41. What two particles interact in Compton scattering?
 - a. photon and electron
 - b. proton and electron
 - c. neutron and electron
 - d. proton and neutron
- **42**. What is the momentum of a 500-nm photon?
 - a. $8.35 \times 10^{-26} \text{ kg} \cdot \text{m/s}$
 - b. $3.31 \times 10^{-40} \text{ kg} \cdot \text{m/s}$
 - c. $7.55 \times 10^{26} \text{ kg} \cdot \text{m/s}$
 - d. $1.33 \times 10^{-27} \text{ kg} \cdot \text{m/s}$
- **43.** The conservation of what fundamental physics principle is behind the technology of solar sails?
 - a. charge
 - b. mass
 - c. momentum
 - d. angular momentum
- **44**. Terms like frequency, amplitude, and period are tied to what component of wave-particle duality?
 - a. neither the particle nor the wave model of light
 - b. both the particle and wave models of light
 - c. the particle model of light
 - d. the wave model of light
- **45**. Why was it beneficial for Compton to scatter electrons using X-rays and not another region of light like microwaves?
 - a. because X-rays are more penetrating than microwaves
 - b. because X-rays have lower frequency than microwaves
 - c. because microwaves have shorter wavelengths than X-rays
 - d. because X-rays have shorter wavelength than microwaves

elliptical path.

- c. The blackbody radiation curve would look like a vertical line.
- d. The blackbody radiation curve would look like a horizontal line.
- **47**. Because there are more gradations to high frequency radiation than low frequency radiation, scientists also thought it possible that a curve titled the *ultraviolet catastrophe* would occur. Explain what the blackbody radiation curve would look like if this were the case.

- a. The curve would steadily increase in intensity with increasing frequency.
- b. The curve would steadily decrease in intensity with increasing frequency.
- c. The curve would be much steeper than in the blackbody radiation graph.
- d. The curve would be much flatter than in the blackbody radiation graph.
- **48**. Energy provided by a light exists in the following quantities: 150 J, 225 J, 300 J. Define one possible quantum of energy and provide an energy state that cannot exist with this quantum.
 - a. 65 J; 450 J cannot exist
 - b. 70 J; 450 J cannot exist
 - c. 75 J; 375 J cannot exist
 - d. 75 J; 100 J cannot exist
- **49**. Why is Planck's recognition of quantum particles considered the dividing line between classical and modern physics?
 - a. Planck recognized that energy is quantized, which was in sync with the classical physics concepts but not in agreement with modern physics concepts.
 - b. Planck recognized that energy is quantized, which was in sync with modern physics concepts but not in agreement with classical physics concepts.
 - c. Prior to Planck's hypothesis, all the classical physics calculations were valid for subatomic particles, but quantum physics calculations were not valid.
 - d. Prior to Planck's hypothesis, all the classical physics calculations were not valid for macroscopic particles, but quantum physics calculations were valid.
- **50.** How many 500-mm microwave photons are needed to supply the 8 kJ of energy necessary to heat a cup of water by 10 degrees Celsius?
 - a. 8.05×10^{28} photons
 - b. 8.05×10^{26} photons
 - c. 2.01×10^{26} photons
 - d. 2.01×10^{28} photons
- **51**. What is the efficiency of a 100-W, 550-nm lightbulb if a photometer finds that 1 × 10²⁰ photons are emitted each second?
 - a. 101 percent
 - b. 72 percent
 - c. 18 percent
 - d. 36 percent
- 52. Rank the following regions of the electromagnetic spectrum by the amount of energy provided per photon: gamma, infrared, microwave, ultraviolet, radio, visible, X-ray.

- a. radio, microwave, infrared, visible, ultraviolet, X-ray, gamma
- b. radio, infrared, microwave, ultraviolet, visible, X-ray, gamma
- c. radio, visible, microwave, infrared, ultraviolet, X-ray, gamma
- d. radio, microwave, infrared, visible, ultraviolet, gamma, X-ray
- **53**. Why are photons of gamma rays and X-rays able to penetrate objects more successfully than ultraviolet radiation?
 - a. Photons of gamma rays and X-rays carry with them less energy.
 - b. Photons of gamma rays and X-rays have longer wavelengths.
 - c. Photons of gamma rays and X-rays have lower frequencies.
 - d. Photons of gamma rays and X-rays carry with them more energy.

21.2 Einstein and the Photoelectric Effect

- **54.** According to wave theory, what is necessary to eject electrons from a surface?
 - a. Enough energy to overcome the binding energy of the electrons at the surface
 - b. A frequency that is higher than that of the electrons at the surface
 - c. Energy that is lower than the binding energy of the electrons at the surface
 - d. A very small number of photons
- **55**. What is the wavelength of EM radiation that ejects 2.00-eV electrons from calcium metal, given that the binding energy is 2.71 eV?
 - a. $16.1 \times 10^5 \text{ m}$
 - b. 6.21×10^{-5} m
 - c. 9.94×10^{-26} m
 - d. 2.63 × 10⁻⁷ m
- 56. Find the wavelength of photons that eject 0.100-eV electrons from potassium, given that the binding energy is 2.24 eV.
 - a. 6.22×10^{-7} m
 - b. 5.92×10^{-5} m
 - c. 1.24×10^{-5} m
 - d. 5.31×10^{-7} m
- 57. How do solar cells utilize the photoelectric effect?
 - a. A solar cell converts all photons that it absorbs to electrical energy using the photoelectric effect.
 - b. A solar cell converts all electrons that it absorbs to electrical energy using the photoelectric effect.
 - c. A solar cell absorbs the photons with energy less

than the energy gap of the material of the solar cell and converts it to electrical energy using the photoelectric effect.

- d. A solar cell absorbs the photons with energy greater than the energy gap of the material of the solar cell and converts it to electrical energy using the photoelectric effect.
- **58.** Explain the advantages of the photoelectric effect to other forms of energy transformation.
 - a. The photoelectric effect is able to work on the Sun's natural energy.
 - b. The photoelectric effect is able to work on energy generated by burning fossil fuels.
 - c. The photoelectric effect can convert heat energy into electrical energy.
 - d. The photoelectric effect can convert electrical energy into light energy.

21.3 The Dual Nature of Light

- **59.** Upon collision, what happens to the frequency of a photon?
 - a. The frequency of the photon will drop to zero.
 - b. The frequency of the photon will remain the same.
 - c. The frequency of the photon will increase.
 - d. The frequency of the photon will decrease.
- **60**. How does the momentum of a photon compare to the momentum of an electron of identical energy?
 - a. Momentum of the photon is greater than the momentum of an electron.
 - b. Momentum of the photon is less than the momentum of an electron.
 - c. Momentum of the photon is equal to the momentum of an electron.
 - d. Momentum of the photon is zero due to zero rest mass but the momentum of an electron is finite.
- **61**. A 500-nm photon strikes an electron and loses 20 percent of its energy. What is the new momentum of the photon?
 - a. $4.24 \times 10^{-27} \text{ kg} \cdot \text{m/s}$
 - b. $3.18 \times 10^{-27} \text{ kg} \cdot \text{m/s}$
 - c. $2.12 \times 10^{-27} \text{ kg} \cdot \text{m/s}$
 - d. $1.06 \times 10^{-27} \text{ kg} \cdot \text{m/s}$
- **62**. A 500-nm photon strikes an electron and loses 20 percent of its energy. What is the speed of the recoiling electron?
 - a. $7.18 \times 10^5 \text{ m/s}$
 - b. $6.18 \times 10^5 \text{ m/s}$

- c. $5.18 \times 10^5 \text{ m/s}$
- d. 4.18 × 10⁵ m/s
- **63**. When a photon strikes a solar sail, what is the direction of impulse on the photon?
 - a. parallel to the sail
 - b. perpendicular to the sail
 - c. tangential to the sail
 - d. opposite to the sail
- **64**. What is a fundamental difference between solar sails and sails that are used on sailboats?
 - Solar sails rely on disorganized strikes from light particles, while sailboats rely on disorganized strikes from air particles.
 - Solar sails rely on disorganized strikes from air particles, while sailboats rely on disorganized strikes from light particles.
 - c. Solar sails rely on organized strikes from air particles, while sailboats rely on organized strikes from light particles.
 - d. Solar sails rely on organized strikes from light particles, while sailboats rely on organized strikes from air particles.
- **65**. The wavelength of a particle is called the de Broglie wavelength, and it can be found with the equation $p = \frac{h}{4}$.

Yes or no—Can the wavelength of an electron match that of a proton?

- a. Yes, a slow-moving electron can achieve the same momentum as a slow-moving proton.
- b. No, a fast-moving electron cannot achieve the same momentum, and hence the same wavelength, as a proton.
- c. No, an electron can achieve the same momentum, and hence not the same wavelength, as a proton.
- d. Yes, a fast-moving electron can achieve the same momentum, and hence have the same wavelength, as a slow-moving proton.
- **66**. Large objects can move with great momentum. Why then is it difficult to see their wave-like nature?
 - a. Their wavelength is equal to the object's size.
 - b. Their wavelength is very small compared to the object's size.
 - c. Their wavelength is very large compared to the object's size.
 - d. Their frequency is very small compared to the object's size.

Extended Response

21.1 Planck and Quantum Nature of Light

- **67**. Some television tubes are CRTs. They use an approximately 30-kV accelerating potential to send electrons to the screen, where the electrons stimulate phosphors to emit the light that forms the pictures we watch. Would you expect X-rays also to be created? Explain.
 - a. No, because the full spectrum of EM radiation is not emitted at any temperature.
 - b. No, because the full spectrum of EM radiation is not emitted at certain temperatures.
 - c. Yes, because the full spectrum of EM radiation is emitted at any temperature.
 - d. Yes, because the full spectrum of EM radiation is emitted at certain temperatures.
- **68**. If Planck's constant were large, say 10³⁴ times greater than it is, we would observe macroscopic entities to be quantized. Describe the motion of a child's swing under such circumstances.
 - a. The child would not be able to swing with particular energies.
 - b. The child could be released from any height.
 - c. The child would be able to swing with constant velocity.
 - d. The child could be released only from particular heights.
- **69**. What is the accelerating voltage of an X-ray tube that produces X-rays with the shortest wavelength of 0.0103 nm?
 - a. $1.21 \times 10^{10} \text{ V}$
 - b. $2.4 \times 10^5 \text{ V}$
 - c. $3.0 \times 10^{-33} \text{ V}$
 - d. $1.21 \times 10^5 \text{ V}$
- **70.** Patients in a doctor's office are rightly concerned about receiving a chest X-ray. Yet visible light is also a form of electromagnetic radiation and they show little concern about sitting under the bright lights of the waiting room. Explain this discrepancy.
 - a. X-ray photons carry considerably more energy so they can harm the patients.
 - b. X-ray photons carry considerably less energy so they can harm the patients.
 - c. X-ray photons have considerably longer wavelengths so they cannot harm the patients.
 - d. X-ray photons have considerably lower frequencies so they can harm the patients.

21.2 Einstein and the Photoelectric Effect

71. When increasing the intensity of light shining on a

metallic surface, it is possible to increase the current created on that surface. Classical theorists would argue that this is evidence that intensity causes charge to move with a greater kinetic energy. Argue this logic from the perspective of a modern physicist.

- a. The increased intensity increases the number of ejected electrons. The increased current is due to the increase in the number of electrons.
- b. The increased intensity decreases the number of ejected electrons. The increased current is due to the decrease in the number of electrons ejected.
- c. The increased intensity does not alter the number of electrons ejected. The increased current is due to the increase in the kinetic energy of electrons.
- d. The increased intensity alters the number of electrons ejected, but an increase in the current is due to an increase in the kinetic energy of electrons.
- **72.** What impact does the quantum nature of electromagnetic radiation have on the understanding of speed at the particle scale?
 - a. Speed must also be quantized at the particle scale.
 - b. Speed will not be quantized at the particle scale.
 - c. Speed must be zero at the particle scale.
 - d. Speed will be infinite at the particle scale.
- **73.** A 500 nm photon of light strikes a semi-conductive surface with a binding energy of 2 eV. With what velocity will an electron be emitted from the semi-conductive surface?
 - a. $8.38 \times 10^5 \text{ m/s}$
 - b. $9.33 \times 10^5 \text{ m/s}$
 - c. $3 \times 10^8 \text{ m/s}$
 - d. 4.11 × 10⁵ m/s
- **74.** True or false—Treating food with ionizing radiation helps keep it from spoiling.
 - a. true
 - b. false

21.3 The Dual Nature of Light

- **75.** When testing atomic bombs, scientists at Los Alamos recognized that huge releases of energy resulted in problems with power and communications systems in the area surrounding the blast site. Explain the possible tie to Compton scattering.
 - a. The release of light energy caused large-scale emission of electrons.
 - b. The release of light energy caused large-scale emission of protons.
 - c. The release of light energy caused large-scale emission of neutrons.
 - d. The release of light energy caused large-scale

emission of photons.

- **76.** Sunlight above the Earth's atmosphere has an intensity of 1.30 kW/m2 . If this is reflected straight back from a mirror that has only a small recoil, the light's momentum is exactly reversed, giving the mirror twice the incident momentum. If the mirror were attached to a solar sail craft, how fast would the craft be moving after 24 hr? Note—The average mass per square meter of the craft is 0.100 kg.
 - a. $8.67 \times 10^{-5} \text{ m/s}^2$
 - b. $8.67 \times 10^{-6} \text{ m/s}^2$
 - c. 94.2 m/s
 - d. 7.49 m/s
- **77.** Consider the counter-clockwise motion of LightSail-1 around Earth. When will the satellite move the fastest?



- a. point A
- b. point B
- c. point C
- d. point D
- **78.** What will happen to the interference pattern created by electrons when their velocities are increased?
 - a. There will be more zones of constructive interference and fewer zones of destructive interference.
 - b. There will be more zones of destructive interference and fewer zones of constructive interference.
 - c. There will be more zones of constructive and destructive interference.
 - d. There will be fewer zones of constructive and destructive interference.